

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



09/102,037

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A23L 1/305	A2	(11) International Publication Number: WO 94/03075 (43) International Publication Date: 17 February 1994 (17.02.94)												
(21) International Application Number: PCT/US93/07190 (22) International Filing Date: 29 July 1993 (29.07.93) (30) Priority data: <table border="0"><tr><td>923,780</td><td>31 July 1992 (31.07.92)</td><td>US</td></tr><tr><td>946,235</td><td>16 September 1992 (16.09.92)</td><td>US</td></tr><tr><td>029,335</td><td>4 March 1993 (04.03.93)</td><td>US</td></tr><tr><td>040,510</td><td>31 March 1993 (31.03.93)</td><td>US</td></tr></table> (71) Applicant: CREATIVE BIOMOLECULES, INC. [US/US]; 45 South Street, Hopkinton, MA 01748 (US). (72) Inventors: KUBERASAMPATH, Thangavel ; Six Spring Street, Medway, MA 02053 (US). COHEN, Charles, M. ; 98 Winthrop Street, Medway, MA 02053 (US). RUEGER, David, C. ; 19 Downey Street, Hopkinton, MA 01748 (US). OPPERMANN, Hermann ; 25 Summer Hill Road, Medway, MA 02053 (US). PANG, Roy, H., L. ; 15 Partridge Road, Etna, NH 03750 (US).		923,780	31 July 1992 (31.07.92)	US	946,235	16 September 1992 (16.09.92)	US	029,335	4 March 1993 (04.03.93)	US	040,510	31 March 1993 (31.03.93)	US	(74) Agent: KELLEY, Robin, D.; Testa, Hurwitz & Thibault, Exchange Place, 53 State Street, Boston, MA 02109 (US). (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
923,780	31 July 1992 (31.07.92)	US												
946,235	16 September 1992 (16.09.92)	US												
029,335	4 March 1993 (04.03.93)	US												
040,510	31 March 1993 (31.03.93)	US												
(54) Title: MORPHOGEN-ENRICHED DIETARY COMPOSITION (57) Abstract Disclosed are methods and compositions useful in dietary applications and capable of enhancing tissue morphogenesis, including tissue development and viability in a mammal, particularly a human. The methods and compositions include a morphogen which, when provided to an individual as a food formulation or supplement, is capable of enhancing tissue development and viability in the individual.														

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

MORPHOGEN-ENRICHED DIETARY COMPOSITION

Field of the Invention

This invention relates generally to the field of
5 dietary compositions and supplements.

Background of the Invention

The present invention relates to compositions
10 useful as mammalian dietary compositions and
supplements. In particular, the invention relates to
food additives and dietary supplements capable of
enhancing tissue morphogenesis and development,
particularly in individuals at risk for normal tissue
15 development and viability. Examples of such
individuals include infants, particularly prematurely-
born ("preterm") and low birth weight infants, and
juveniles; aged individuals; and individuals
experiencing altered metabolic function and/or
20 suffering from metabolic dysfunctions and other
disorders that threaten organ or tissue function or
viability, such as can result from malnutrition or
starvation, autoimmune diseases, organ cirrhosis and
other tissue necrotizing dysfunctions, or disorders
25 associated with aging cells (cell senescence.)

Mammalian infants are nourished by mother's milk
until such time as they can digest food solids. Infant
formulas now exist for humans and other mammals which
30 can supplant or supplement mother's milk. The formulas
may be milk based (e.g., cow milk) or non-milk-based

- 2 -

(e.g., soy). Particularly at risk are prematurely born infants whose tissues and organs are at an earlier stage of development, and whose nutritional requirements may differ from those of full term

5 infants. Formula development is an ongoing endeavor to more accurately mimic the beneficial aspects of mother's milk. Nevertheless, despite the efforts of many researchers, infant formulas still differ in a number of significant ways from human milk. In part

10 this is due because human milk has many substances, such as immunoglobulins, free amino acids, polyamines, nucleotides and polyunsaturated fatty acids not present, for example, in cow's milk. In addition, while infant formulas try to mimic the protein quantity

15 found in human milk, the foreign proteins typically are present in the formula as hydrolysates to avoid rejection or reaction by the infant's digestive system. The proteins are present primarily as amino acid sources rather than as functional proteins as might

20 normally be transmitted by the nursing mother to the infant. In addition, human milk may contain unidentified growth and differentiation factors that are important for overall tissue and skeletal development.

25

Another group of individuals with potentially unique nutritional requirements are individuals undergoing metabolic changes which may result from periods of intense growth or stress, including, for

30 example, pregnant women and drowning victims. Other sources of stress to the body may result from

- 3 -

malnutrition or starvation, or from metabolic disorders that affect organ viability, such as autoimmune disease and organ cirrhosis. Aged individuals, and postmenopausal women also have altered or slower
5 metabolic function. All of these individuals are at risk for tissue damage or loss of tissue function due to altered metabolic function.

Reduced or lost tissue function due to malnutrition
10 also is found in many patients admitted to hospitals (protein energy malnutrition, "PEM"). Proper nutritional support for such patients, while not a primary mode of treatment is, nevertheless, an important factor for therapy and recovery. It is,
15 therefore important to administer a nutritionally balanced diet given orally, enterally or parenterally, adequate to the needs of the patient. This is especially true for those patients where conventional feeding is contraindicated (e.g., in dehydrated or
20 gastroenterological patients) or is insufficient (e.g., in hypercatabolic patients). The enteral or oral mode of administration of foods typically is preferable to parenteral modes because of the lower morbidity, trophic effect upon the intestinal mucosa, reduced
25 dependency on instrumentation and lower costs.

It is an object of this invention to provide dietary compositions and supplements for enhancing tissue morphogenesis, including tissue growth,
30 development, maintenance and viability in a mammal, particularly a human. Another object of the invention is to provide an infant formula capable of enhancing tissue development in an infant or juvenile. Still another object is to provide an an infant formula that

- 4 -

more closely mimics a nursing mother's milk. Another object of the invention is to provide dietary supplements for individuals at risk for normal tissue development, growth, maintenance and viability, including premature infants, aged individuals and individuals with altered metabolic function and/or suffering from disorders or metabolic dysfunctions which threaten organ viability and function. These and other objects and features of the invention will be apparent from the description, drawings, and claims which follow.

Summary of the Invention

The present invention provides compositions and methods useful in dietary applications and capable of enhancing tissue morphogenesis, including tissue growth, development, maintenance and viability in a mammal, particularly a human. The dietary compositions and supplements of this invention comprise a morphogenic protein ("morphogen"), as described herein, which, when provided to an individual as a food formulation or supplement, is capable of enhancing tissue development, growth, maintenance and/or viability in the individual. The compositions and processes provided herein are suitable for both infants and adults, and as part of clinical nutrition.

As used herein, "enhancing tissue viability" is understood to mean protecting tissue from lost or reduced tissue function due to cell damage or cell senescence, including inducing cells to maintain their differentiated phenotype, inducing regeneration of damaged tissue, and/or inhibiting additional damage

- 5 -

thereto. "Morphogenically effective concentration" is understood to mean a concentration sufficient to enhance tissue development and tissue viability in an individual at risk for tissue damage and/or reduced or lost tissue function due to insufficient nutritional considerations, tissue damage associated therewith, and/or incomplete tissue development, regardless of etiology. The ability of morphogens to repair, regenerate and protect various disparate tissues, including but not limited to, tissues of the gastrointestinal tract, including the oral mucosa, liver tissue, dentin tissue, periodontal tissue, nerve tissue, bone tissue, and any tissue at risk of damage due to immune response-mediated tissue destruction, including ischemia-reperfusion related tissue damage are disclosed in international applications US 92/01968 (WO 92/15323), US 92/07358 (WO 93/04692) and US 92/07232 (WO 93/05751) respectively, the disclosures of which are incorporated herein by reference.

"Morphogen-solubilizing molecule" is understood to mean a molecule capable of maintaining a morphogen in soluble form in physiologically buffered solutions. "Food formulation" is understood to mean a dietary composition normally ingested by an individual to satisfy the body's fundamental nutritional requirements; "dietary supplement" is understood to mean supplemental compositions ingested by an individual in addition to the food formulations ingested to satisfy the fundamental nutritional requirements. Multivitamin and iron tablets are examples of common dietary supplements. "Dietary composition" is understood to include both food formulations and dietary supplements. As used herein, the term "infant formula" is understood to refer to the

- 6 -

well established infant compositions as defined by the American Academy of Pediatrics (AAP) and the AAP Committee on Nutrition ((1985) Pediatrics 75:976, the European Society of Pediatric Gastroenterology and Nutrition (ESPGAN) and the ESPGAN Committee on Nutrition ((1987) Acta Paed Scan Suppl:330), including recent updates published by these committees on infant formula nutritional guidelines.

- 10 The dietary composition or supplement preferably is administered orally, and may be provided in liquid form or as a powder to be dissolved in a beverage. Alternatively, the dietary supplement may be provided as a solid, e.g., in a capsular, tablet, troche or
15 lozenge form; or, the supplement may be provided as an aerosol, for oral or nasal administration. Where oral administration is not possible or desirable, other administration routes are envisioned. For example, for some premature infants, or for intubated patients,
20 parenteral administration may be required, e.g., via an enteral feeding tube.

- The morphogen may be provided alone or in association with one or more suitable excipients or
25 carriers, and/or in combination with other beneficial molecules such as vitamins, minerals, lipids, fiber sources and the like. The dietary supplements also may include pharmaceutically acceptable inert materials for use as binders or stabilizers, including magnesium
30 stearate or calcium carbonate. The morphogen may be

- 7 -

formulated together with one or more normal food ingredients, e.g., as part of a food formulation. Alternatively or, in addition, the morphogen may be provided as a dietary supplement in, for example, 5 tablet or syrup form.

The mature form of the morphogen, or active truncated forms thereof which may be formulated in the composition, further may be provided in association 10 with a morphogen precursor "pro" domain, which is known to enhance the solubility of the protein in physiologically buffered solutions. Other useful molecules known to enhance protein solubility include casein, including derivatives, salts and analogs 15 thereof, as well as other milk components, and various serum and milk serum proteins. Additional useful molecules which may be associated with the morphogen include tissue targeting molecules capable of directing the morphogen to a desired target tissue. Tissue 20 targeting molecules envisioned to be useful in the treatment protocols of this invention include antibodies, antibody fragments or other binding proteins which interact specifically with surface molecules on the target tissue cells.

25

Still another useful tissue targeting molecule may be part or all of a morphogen precursor "pro" domain. Morphogens may be synthesized in one tissue and secreted and transported to another tissue. For 30 example, while the protein has been shown to be active in bone tissue, the primary source of OP-1 synthesis appears to be the tissue of the urogenic system (e.g., renal and bladder tissue), with secondary expression levels occurring in the brain, heart, lungs and

- 8 -

gastrointestinal tract (GI tract, see below.)
Moreover, the protein has been identified in serum,
saliva and various milk forms. In addition, the
secreted form of the protein comprises the mature dimer
5 in association with the pro domain of the intact
morphogen sequence. Accordingly, the associated
morphogen pro domains may act to target specific
morphogens to different tissues in vivo. As described
below, morphogen species comprising the pro domain may
10 be obtained from the culture medium of morphogen-
secreting mammalian cells. Alternatively, a tissue-
targeting species may be formulated by complexing the
mature dimer (or an active fragment thereof) with part
or all of a pro domain.

15

Associated tissue targeting or solubility-enhancing
molecules also may be covalently linked to the
morphogen using standard chemical means.

20 In one preferred embodiment, the morphogen
comprises part of an infant formula. The infant formula
may be milk-based or nonmilk-based, e.g., soy-based. A
typical ready-to-feed morphogen-enriched formulation
for infants, when diluted to feeding concentrations,
25 comprises, in addition to the morphogen added to the
formula, from about 1-5% by weight fat, from about 0.01
to about 0.5% by weight immunoglobulins as appropriate,
from about 4-10% by weight carbohydrate in a quantity
substantially to mimic the carbohydrate content of
30 human mother's milk, from about 0.5 to 4% by weight

- 9 -

protein in a quantity substantially to mimic the protein content of human mother's milk, optional vitamins and minerals as required, a total solids content of from about 8 to 17% by weight, and the
5 remainder water.

In another preferred embodiment, the dietary composition is formulated for individuals at risk for reduced or lost tissue function, such as postmenopausal
10 women, elderly individuals, undernourished or malnourished individuals, dehydrated individuals, drowning victims, individuals suffering from metabolic disorders including an endocrine imbalance, gastrointestinal disorders, or immune-compromised
15 individuals. Undernourished or malnourished individuals include those suffering from a lack of food (starvation) and/or eating disorders (e.g., anorexia nervosa), and/or suffering from a malabsorption syndrome (e.g., individuals afflicted with digestive or
20 intestinal fistulas, shortened bowel, or hypercatabolism.) Individuals receiving a medical therapy, including radiotherapy, chemotherapy or a surgical procedure also are at risk for reduced or lost tissue function as a result of a therapy-related
25 malabsorption-malnutrition dysfunction. In another embodiment, the dietary supplement is formulated for individuals undergoing periods of increased growth or stress, such as infants and juveniles, or pregnant or lactating women. In another embodiment, the dietary
30 supplement is formulated for individuals at risk for reduced or lost organ function as results from tissue cirrhosis or an autoimmune disease.

- 10 -

Morphogen-enriched nutritional products, particularly clinical nutrition products for use in hospital or other clinical settings, in addition to comprising a morphogen preferably are based on the utilization of diverse other protein sources (casein, sodium and calcium caseinate, isolated soy protein, protein hydrolyzates and/or crystalline amino acids) mixtures of vegetable and animal fats, carbohydrates (basically glucose polymers), vitamins and minerals to meet, at least, the dietary intakes recommended for healthy individuals (see, for example, Committee on Dietary Allowances, Food and Nutrition Board, Nat Acad Sci, 9th Ed, 1980).

Among the morphogens useful in this invention are proteins originally identified as osteogenic proteins, such as the OP-1, OP-2 and CBMP2 proteins, as well as amino acid sequence-related proteins such as DPP (from *Drosophila*), Vgl (from *Xenopus*), Vgr-1 (from mouse, see U.S. 5,011,691 to Oppermann et al.), GDF-1 (from mouse, see Lee (1991) PNAS 88:4250-4254), all of which are presented in Table II and Seq. ID Nos.5-14), and the recently identified 60A protein (from *Drosophila*, Seq. ID No. 24, see Wharton et al. (1991) PNAS 88:9214-9218.) The members of this family, which include members of the TGF- β super-family of proteins, share substantial amino acid sequence homology in their C-terminal regions. The proteins are translated as a precursor, having an N-terminal signal peptide sequence, typically less than about 30 residues, followed by a "pro" domain that is cleaved to yield the mature sequence. The "pro" form of the protein includes the pro domain and the mature domain, and forms a soluble species that appears to be the primary

- 11 -

form secreted from cultured mammalian cells. The signal peptide is cleaved rapidly upon translation, at a cleavage site that can be predicted in a given sequence using the method of Von Heijne ((1986) Nucleic
 5 Acids Research 14:4683-4691.) Table I, below, describes the various morphogens identified to date, including their nomenclature as used herein, their Seq. ID references, and publication sources for the amino acid sequences for the full length proteins not
 10 included in the Seq. Listing. The disclosure of these publications is incorporated herein by reference.

TABLE I

15	"OP-1"	Refers generically to the group of morphogenically active proteins expressed from part or all of a DNA sequence encoding OP-1 protein, including allelic and species variants thereof, e.g., human
20		OP-1 ("hOP-1", Seq. ID No. 5, mature protein amino acid sequence), or mouse OP-1 ("mOP-1", Seq. ID No. 6, mature protein amino acid sequence.) The
25		conserved seven cysteine skeleton is defined by residues 38 to 139 of Seq. ID Nos. 5 and 6. The cDNA sequences and the amino acids encoding the full length
30		proteins are provided in Seq. ID Nos. 16 and 17 (hOP1) and Seq. ID Nos. 18 and 19 (mOP1.) The mature proteins are defined

- 12 -

by residues 293-431 (hOP1) and 292-430 (mOP1). The "pro" regions of the proteins, cleaved to yield the mature, morphogenically active proteins are defined essentially by residues 30-292 (hOP1) and residues 30-291 (mOP1).

"OP-2" refers generically to the group of active proteins expressed from part or all of a DNA sequence encoding OP-2 protein, including allelic and species variants thereof, e.g., human OP-2 ("hOP-2", Seq. ID No. 7, mature protein amino acid sequence) or mouse OP-2 ("mOP-2", Seq. ID No. 8, mature protein amino acid sequence). The conserved seven cysteine skeleton is defined by residues 38 to 139 of Seq. ID Nos. 7 and 8. The cDNA sequences and the amino acids encoding the full length proteins are provided in Seq. ID Nos. 20 and 21 (hOP2) and Seq. ID Nos. 22 and 23 (mOP2.) The mature proteins are defined essentially by residues 264-402 (hOP2) and 261-399 (mOP2). The "pro" regions of the proteins, cleaved to yield the mature, morphogenically active proteins likely are defined essentially by residues 18-263 (hOP2) and residues 18-260 (mOP2). (Another cleavage site also occurs 21 residues upstream for both OP-2 proteins.)

- 13 -

- "CBMP2" refers generically to the morphogenically active proteins expressed from a DNA sequence encoding the CBMP2 proteins, including allelic and species variants thereof, e.g., human CBMP2A ("CBMP2A(fx)", Seq ID No. 9) or human CBMP2B DNA ("CBMP2B(fx)", Seq. ID No. 10). The amino acid sequence for the full length proteins, referred to in the literature as BMP2A and BMP2B, or BMP2 and BMP4, appear in Wozney, et al. (1988) Science 242:1528-1534. The pro domain for BMP2 (BMP2A) likely includes residues 25-248 or 25-282; the mature protein, residues 249-396 or 283-396. The pro domain for BMP4 (BMP2B) likely includes residues 25-256 or 25-292; the mature protein, residues 257-408 or 293-408.
- 20 "DPP(fx)" refers to protein sequences encoded by the *Drosophila* DPP gene and defining the conserved seven cysteine skeleton (Seq. ID No. 11). The amino acid sequence for the full length protein appears in Padgett, et al (1987) Nature 325: 81-84. The pro domain likely extends from the signal peptide cleavage site to residue 456; the mature protein likely is defined by residues 457-588.
- 30 "Vgl(fx)" refers to protein sequences encoded by the *Xenopus* Vgl gene and defining the conserved seven cysteine skeleton (Seq. ID No. 12). The amino acid sequence for the

- 14 -

5 full length protein appears in Weeks (1987) Cell 51: 861-867. The prodomain likely extends from the signal peptide cleavage site to residue 246; the mature protein likely is defined by residues 247-360.

10 "Vgr-1(fx)" refers to protein sequences encoded by the murine Vgr-1 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 13). The amino acid sequence for the full length protein appears in Lyons, et al, (1989) PNAS 86: 4554-4558. The prodomain likely extends from the signal peptide cleavage site to residue 299; the mature protein likely is defined by residues 300-438.

20 "GDF-1(fx)" refers to protein sequences encoded by the human GDF-1 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 14). The cDNA and encoded amino sequence for the full length protein is provided in Seq. ID. No. 32. The prodomain likely extends from the signal peptide clavage site to residue 214; the mature protein likely is defined by residues 215-372.

30 "60A" refers generically to the morphogenically active proteins expressed from part or all of a DNA sequence (from the Drosophila 60A gene) encoding the 60A proteins (see Seq. ID No. 24 wherein the cDNA and encoded

- 15 -

amino acid sequence for the full length protein is provided). "60A(fx)" refers to the protein sequences defining the conserved seven cysteine skeleton
5 (residues 354 to 455 of Seq. ID No. 24.)
The prodomain likely extends from the signal peptide cleavage site to residue 324; the mature protein likely is defined by residues 325-455.

10 "BMP3(fx)" refers to protein sequences encoded by the human BMP3 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 26).
The amino acid sequence for the full
15 length protein appears in Wozney et al. (1988) Science 242: 1528-1534. The pro domain likely extends from the signal peptide cleavage site to residue 290; the mature protein likely is defined by
20 residues 291-472.

"BMP5(fx)" refers to protein sequences encoded by the human BMP5 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 27).
25 The amino acid sequence for the full length protein appears in Celeste, et al. (1991) PNAS 87: 9843-9847. The pro domain likely extends from the signal peptide cleavage site to residue 316; the mature
30 protein likely is defined by residues 317-454.

- 16 -

"BMP6(fx)" refers to protein sequences encoded by the human BMP6 gene and defining the conserved seven cysteine skeleton (Seq. ID No. 28). The amino acid sequence for the full length protein appears in Celeste, et al. (1990) PNAS 87: 9843-5847. The pro domain likely includes extends from the signal peptide cleavage site to residue 374; the mature sequence likely includes residues 375-513.

The OP-2 proteins have an additional cysteine residue in this region (e.g., see residue 41 of Seq. ID Nos. 7 and 8), in addition to the conserved cysteine skeleton in common with the other proteins in this family. The GDF-1 protein has a four amino acid insert within the conserved skeleton (residues 44-47 of Seq. ID No. 14) but this insert likely does not interfere with the relationship of the cysteines in the folded structure. In addition, the CBMP2 proteins are missing one amino acid residue within the cysteine skeleton.

The morphogens are inactive when reduced, but are active as oxidized homodimers and when oxidized in combination with other morphogens of this invention. Thus, as defined herein, a morphogen is a dimeric protein comprising a pair of polypeptide chains, wherein each polypeptide chain comprises at least the C-terminal six cysteine skeleton defined by residues 43-139 of Seq. ID No. 5, including functionally equivalent arrangements of these cysteines (e.g., amino acid insertions or deletions which alter the linear arrangement of the cysteines in the sequence but not

- 17 -

their relationship in the folded structure), such that, when the polypeptide chains are folded, the dimeric protein species comprising the pair of polypeptide chains has the appropriate three-dimensional structure, including the appropriate intra- or inter-chain disulfide bonds such that the protein is capable of acting as a morphogen as defined herein. Specifically, the morphogens generally are capable of all of the following biological functions in a morphogenically permissive environment: stimulating proliferation of progenitor cells; stimulating the differentiation of progenitor cells; stimulating the proliferation of differentiated cells, and supporting the growth and maintenance of differentiated cells. In addition, it is also anticipated that these morphogens are capable of inducing redifferentiation of committed cells under appropriate environmental conditions.

In one preferred aspect, the morphogens of this invention comprise one of two species of generic amino acid sequences: Generic Sequence 1 (Seq. ID No. 1) or Generic Sequence 2 (Seq. ID No. 2); where each Xaa indicates one of the 20 naturally-occurring L-isomer, α -amino acids or a derivative thereof. Generic Sequence 1 comprises the conserved six cysteine skeleton and Generic Sequence 2 comprises the conserved six cysteine skeleton plus the additional cysteine identified in OP-2 (see residue 36, Seq. ID No. 2). In another preferred aspect, these sequences further comprise the following additional sequence at their N-terminus:

Cys Xaa Xaa Xaa Xaa (Seq. ID No. 15)

1

5

- 18 -

Preferred amino acid sequences within the foregoing generic sequences include: Generic Sequence 3 (Seq. ID No. 3), Generic Sequence 4 (Seq. ID No. 4), Generic Sequence 5 (Seq. ID No. 30) and Generic Sequence 6 (Seq. ID No. 31), listed below. These Generic Sequences accommodate the homologies shared among the various preferred members of this morphogen family identified in Table II, as well as the amino acid sequence variation among them. Specifically, Generic Sequences 3 and 4 are composite amino acid sequences of the following proteins presented in Table II and identified in Seq. ID Nos. 5-14: human OP-1 (hOP-1, Seq. ID Nos. 5 and 16-17), mouse OP-1 (mOP-1, Seq. ID Nos. 6 and 18-19), human and mouse OP-2 (Seq. ID Nos. 7, 8, and 20-22), CBMP2A (Seq. ID No. 9), CBMP2B (Seq. ID No. 10), DPP (from *Drosophila*, Seq. ID No. 11), Vgl, (from *Xenopus*, Seq. ID No. 12), Vgr-1 (from mouse, Seq. ID No. 13), and GDF-1 (from mouse, Seq. ID No. 14.) The generic sequences include both the amino acid identity shared by the sequences in Table II, as well as alternative residues for the variable positions within the sequence. Note that these generic sequences allow for an additional cysteine at position 41 or 46 in Generic Sequences 3 or 4, respectively, providing an appropriate cysteine skeleton where inter- or intramolecular disulfide bonds can form, and contain certain critical amino acids which influence the tertiary structure of the proteins.

- 19 -

Generic Sequence 3

Leu Tyr Val Xaa Phe

	1		5
	Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa		
5			10
	Xaa Ala Pro Xaa Gly Xaa Xaa Ala		
	15		20
	Xaa Tyr Cys Xaa Gly Xaa Cys Xaa		
	25		30
10	Xaa Pro Xaa Xaa Xaa Xaa Xaa		
			35
	Xaa Xaa Xaa Asn His Ala Xaa Xaa		
	40		45
	Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa		
15			50
	Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys		
	55		60
	Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa		
			65
20	Xaa Xaa Xaa Leu Xaa Xaa Xaa		
	70		75
	Xaa Xaa Xaa Xaa Val Xaa Leu Xaa		
			80
	Xaa Xaa Xaa Xaa Met Xaa Val Xaa		
25	85		90

- 20 -

Xaa Cys Gly Cys Xaa

95

wherein each Xaa is independently selected from a group of one or more specified amino acids defined as follows: "Res." means "residue" and Xaa at res.4 = (Ser, Asp or Glu); Xaa at res.6 = (Arg, Gln, Ser or Lys); Xaa at res.7 = (Asp or Glu); Xaa at res.8 = (Leu or Val); Xaa at res.11 = (Gln, Leu, Asp, His or Asn); Xaa at res.12 = (Asp, Arg or Asn); Xaa at res.14 = (Ile or Val); Xaa at res.15 = (Ile or Val); Xaa at res.18 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.20 = (Tyr or Phe); Xaa at res.21 = (Ala, Ser, Asp, Met, His, Leu or Gln); Xaa at res.23 = (Tyr, Asn or Phe); Xaa at res.26 = (Glu, His, Tyr, Asp or Gln); Xaa at res.28 = (Glu, Lys, Asp or Gln); Xaa at res.30 = (Ala, Ser, Pro or Gln); Xaa at res.31 = (Phe, Leu or Tyr); Xaa at res.33 = (Leu or Val); Xaa at res.34 = (Asn, Asp, Ala or Thr); Xaa at res.35 = (Ser, Asp, Glu, Leu or Ala); Xaa at res.36 = (Tyr, Cys, His, Ser or Ile); Xaa at res.37 = (Met, Phe, Gly or Leu); Xaa at res.38 = (Asn or Ser); Xaa at res.39 = (Ala, Ser or Gly); Xaa at res.40 = (Thr, Leu or Ser); Xaa at res.44 = (Ile or Val); Xaa at res.45 = (Val or Leu); Xaa at res.46 = (Gln or Arg); Xaa at res.47 = (Thr, Ala or Ser); Xaa at res.49 = (Val or Met); Xaa at res.50 = (His or Asn); Xaa at res.51 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.52 = (Ile, Met, Asn, Ala or Val); Xaa at res.53 = (Asn, Lys, Ala or Glu); Xaa at res.54 = (Pro or Ser); Xaa at res.55 = (Glu, Asp, Asn, or Gly); Xaa at res.56 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser or Ala); Xaa at res.57 = (Val, Ala or Ile); Xaa at res.58 = (Pro or Asp); Xaa at res.59 = (Lys or Leu); Xaa at res.60 = (Pro or Ala); Xaa at res.63 = (Ala or Val); Xaa at

- 21 -

res.65 = (Thr or Ala); Xaa at res.66 = (Gln, Lys, Arg or Glu); Xaa at res.67 = (Leu, Met or Val); Xaa at res.68 = (Asn, Ser or Asp); Xaa at res.69 = (Ala, Pro or Ser); Xaa at res.70 = (Ile, Thr or Val); Xaa at
5 res.71 = (Ser or Ala); Xaa at res.72 = (Val or Met); Xaa at res.74 = (Tyr or Phe); Xaa at res.75 = (Phe, Tyr or Leu); Xaa at res.76 = (Asp or Asn); Xaa at res.77 = (Asp, Glu, Asn or Ser); Xaa at res.78 = (Ser, Gln, Asn or Tyr); Xaa at res.79 = (Ser, Asn, Asp or Glu); Xaa at
10 res.80 = (Asn, Thr or Lys); Xaa at res.82 = (Ile or Val); Xaa at res.84 = (Lys or Arg); Xaa at res.85 = (Lys, Asn, Gln or His); Xaa at res.86 = (Tyr or His); Xaa at res.87 = (Arg, Gln or Glu); Xaa at res.88 = (Asn, Glu or Asp); Xaa at res.90 = (Val, Thr or Ala);
15 Xaa at res.92 = (Arg, Lys, Val, Asp or Glu); Xaa at res.93 = (Ala, Gly or Glu); and Xaa at res.97 = (His or Arg);

Generic Sequence 4

```

20      Cys Xaa Xaa Xaa Xaa Leu Tyr Val Xaa Phe
        1                               5                               10
      Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
        15
25      Xaa Ala Pro Xaa Gly Xaa Xaa Ala
        20                               25
      Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
        30                               35
      Xaa Pro Xaa Xaa Xaa Xaa Xaa
        40
30

```

- 22 -

```

Xaa Xaa Xaa Asn His Ala Xaa Xaa
      45                      50
Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa
      55
5   Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
      60                      65
    Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
      70
Xaa Xaa Xaa Leu Xaa Xaa Xaa
10   75                      80
    Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
      85
    Xaa Xaa Xaa Xaa Met Xaa Val Xaa
      90                      95
15   Xaa Cys Gly Cys Xaa
      100

```

wherein each Xaa is independently selected from a group of one or more specified amino acids as defined by the following: "Res." means "residue" and Xaa at res.2 =

20 (Lys or Arg); Xaa at res.3 = (Lys or Arg); Xaa at res.4 = (His or Arg); Xaa at res.5 = (Glu, Ser, His, Gly, Arg or Pro); Xaa at res.9 = (Ser, Asp or Glu); Xaa at res.11 = (Arg, Gln, Ser or Lys); Xaa at res.12 = (Asp or Glu); Xaa at res.13 = (Leu or Val); Xaa at res.16 =

25 (Gln, Leu, Asp, His or Asn); Xaa at res.17 = (Asp, Arg, or Asn); Xaa at res.19 = (Ile or Val); Xaa at res.20 = (Ile or Val); Xaa at res.23 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.25 = (Tyr or Phe); Xaa at res.26 = (Ala, Ser, Asp, Met, His, Leu, or Gln); Xaa at res.28 =

30 (Tyr, Asn or Phe); Xaa at res.31 = (Glu, His, Tyr, Asp or Gln); Xaa at res.33 = Glu, Lys, Asp or Gln; Xaa at res.35 = (Ala, Ser or Pro); Xaa at res.36 = (Phe, Leu or Tyr); Xaa at res.38 = (Leu or Val); Xaa at res.39 = (Asn, Asp, Ala or Thr); Xaa at res.40 = (Ser, Asp, Glu,

- 23 -

Leu or Ala); Xaa at res.41 = (Tyr, Cys, His, Ser or Ile); Xaa at res.42 = (Met, Phe, Gly or Leu); Xaa at res.44 = (Ala, Ser or Gly); Xaa at res.45 = (Thr, Leu or Ser); Xaa at res.49 = (Ile or Val); Xaa at res.50 =
5 (Val or Leu); Xaa at res.51 = (Gln or Arg); Xaa at res.52 = (Thr, Ala or Ser); Xaa at res.54 = (Val or Met); Xaa at res.55 = (His or Asn); Xaa at res.56 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.57 = (Ile, Met, Asn, Ala or Val); Xaa at res.58 = (Asn, Lys, Ala
10 or Glu); Xaa at res.59 = (Pro or Ser); Xaa at res.60 = (Glu, Asp, or Gly); Xaa at res.61 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser or Ala); Xaa at res.62 = (Val, Ala or Ile); Xaa at res.63 = (Pro or Asp); Xaa at res.64 = (Lys or Leu); Xaa at res.65 = (Pro or Ala); Xaa at
15 res.68 = (Ala or Val); Xaa at res.70 = (Thr or Ala); Xaa at res.71 = (Gln, Lys, Arg or Glu); Xaa at res.72 = (Leu, Met or Val); Xaa at res.73 = (Asn, Ser or Asp); Xaa at res.74 = (Ala, Pro or Ser); Xaa at res.75 = (Ile, Thr or Val); Xaa at res.76 = (Ser or Ala); Xaa at
20 res.77 = (Val or Met); Xaa at res.79 = (Tyr or Phe); Xaa at res.80 = (Phe, Tyr or Leu); Xaa at res.81 = (Asp or Asn); Xaa at res.82 = (Asp, Glu, Asn or Ser); Xaa at res.83 = (Ser, Gln, Asn or Tyr); Xaa at res.84 = (Ser, Asn, Asp or Glu); Xaa at res.85 = (Asn, Thr or Lys);
25 Xaa at res.87 = (Ile or Val); Xaa at res.89 = (Lys or Arg); Xaa at res.90 = (Lys, Asn, Gln or His); Xaa at res.91 = (Tyr or His); Xaa at res.92 = (Arg, Gln or Glu); Xaa at res.93 = (Asn, Glu or Asp); Xaa at res.95 = (Val, Thr or Ala); Xaa at res.97 = (Arg, Lys, Val,
30 Asp or Glu); Xaa at res.98 = (Ala, Gly or Glu); and Xaa at res.102 = (His or Arg).

- 24 -

Similarly, Generic Sequence 5 (Seq. ID No. 30) and Generic Sequence 6 (Seq. ID No. 31) accommodate the homologies shared among all the morphogen protein family members identified in Table II. Specifically, 5 Generic Sequences 5 and 6 are composite amino acid sequences of human OP-1 (hOP-1, Seq. ID Nos. 5 and 16-17), mouse OP-1 (mOP-1, Seq. ID Nos. 6 and 18-19), human and mouse OP-2 (Seq. ID Nos. 7, 8, and 20-22), CBMP2A (Seq. ID No. 9), CBMP2B (Seq. ID No. 10), DPP 10 (from *Drosophila*, Seq. ID No. 11), Vgl, (from *Xenopus*, Seq. ID No. 12), Vgr-1 (from mouse, Seq. ID No. 13), and GDF-1 (from mouse, Seq. ID No. 14), human BMP3 (Seq. ID No. 26), human BMP5 (Seq. ID No. 27), human BMP6 (Seq. ID No. 28) and 60(A) (from *Drosophila*, Seq. 15 ID Nos. 24-25). The generic sequences include both the amino acid identity shared by these sequences in the C-terminal domain, defined by the six and seven cysteine skeletons (Generic Sequences 5 and 6, respectively), as well as alternative residues for the 20 variable positions within the sequence. As for Generic Sequences 3 and 4, Generic Sequences 5 and 6 allow for an additional cysteine at position 41 (Generic Sequence 5) or position 46 (Generic Sequence 6), providing an appropriate cysteine skeleton where inter- or 25 intramolecular disulfide bonds can form, and containing certain critical amino acids which influence the tertiary structure of the proteins.

- 25 -

Generic Sequence 5

Leu Xaa Xaa Xaa Phe
1 5
5 Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
10
Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala
15 20
Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
10 25 30
Xaa Pro Xaa Xaa Xaa Xaa Xaa
35
Xaa Xaa Xaa Asn His Ala Xaa Xaa
40 45
15 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
50
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
55 60
Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
20 65
Xaa Xaa Xaa Leu Xaa Xaa Xaa
70 75

- 26 -

Xaa Xaa Xaa Xaa Val Xaa Leu Xaa

80

Xaa Xaa Xaa Xaa Met Xaa Val Xaa

85

90

5 Xaa Cys Xaa Cys Xaa

95

wherein each Xaa is independently selected from a group of one or more specified amino acids defined as follows: "Res." means "residue" and Xaa at res.2 =

10 (Tyr or Lys); Xaa at res.3 = Val or Ile); Xaa at res.4 = (Ser, Asp or Glu); Xaa at res.6 = (Arg, Gln, Ser, Lys or Ala); Xaa at res.7 = (Asp, Glu or Lys); Xaa at res.8 = (Leu, Val or Ile); Xaa at res.11 = (Gln, Leu, Asp, His, Asn or Ser); Xaa at res.12 = (Asp, Arg, Asn or

15 Glu); Xaa at res.14 = (Ile or Val); Xaa at res.15 = (Ile or Val); Xaa at res.16 (Ala or Ser); Xaa at res.18 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.19 = (Gly or Ser); Xaa at res.20 = (Tyr or Phe); Xaa at res.21 = (Ala, Ser, Asp, Met, His, Gln, Leu or Gly);

20 Xaa at res.23 = (Tyr, Asn or Phe); Xaa at res.26 = (Glu, His, Tyr, Asp, Gln or Ser); Xaa at res.28 = (Glu, Lys, Asp, Gln or Ala); Xaa at res.30 = (Ala, Ser, Pro, Gln or Asn); Xaa at res.31 = (Phe, Leu or Tyr); Xaa at res.33 = (Leu, Val or Met); Xaa at res.34 = (Asn, Asp,

25 Ala, Thr or Pro); Xaa at res.35 = (Ser, Asp, Glu, Leu, Ala or Lys); Xaa at res.36 = (Tyr, Cys, His, Ser or Ile); Xaa at res.37 = (Met, Phe, Gly or Leu); Xaa at res.38 = (Asn, Ser or Lys); Xaa at res.39 = (Ala, Ser, Gly or Pro); Xaa at res.40 = (Thr, Leu or Ser); Xaa at

30 res.44 = (Ile, Val or Thr); Xaa at res.45 = (Val, Leu or Ile); Xaa at res.46 = (Gln or Arg); Xaa at res.47 = (Thr, Ala or Ser); Xaa at res.48 = (Leu or Ile); Xaa at

- 27 -

res.49 = (Val or Met); Xaa at res.50 = (His, Asn or Arg); Xaa at res.51 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.52 = (Ile, Met, Asn, Ala, Val or Leu); Xaa at res.53 = (Asn, Lys, Ala, Glu, Gly or Phe); Xaa at
5 res.54 = (Pro, Ser or Val); Xaa at res.55 = (Glu, Asp, Asn, Gly, Val or Lys); Xaa at res.56 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser, Ala, Pro or His); Xaa at res.57 = (Val, Ala or Ile); Xaa at res.58 = (Pro or Asp); Xaa at res.59 = (Lys, Leu or Glu); Xaa at res.60 = (Pro or
10 Ala); Xaa at res.63 = (Ala or Val); Xaa at res.65 = (Thr, Ala or Glu); Xaa at res.66 = (Gln, Lys, Arg or Glu); Xaa at res.67 = (Leu, Met or Val); Xaa at res.68 = (Asn, Ser, Asp or Gly); Xaa at res.69 = (Ala, Pro or Ser); Xaa at res.70 = (Ile, Thr, Val or Leu); Xaa at
15 res.71 = (Ser, Ala or Pro); Xaa at res.72 = (Val, Met or Ile); Xaa at res.74 = (Tyr or Phe); Xaa at res.75 = (Phe, Tyr, Leu or His); Xaa at res.76 = (Asp, Asn or Leu); Xaa at res.77 = (Asp, Glu, Asn or Ser); Xaa at res.78 = (Ser, Gln, Asn, Tyr or Asp); Xaa at res.79 =
20 (Ser, Asn, Asp, Glu or Lys); Xaa at res.80 = (Asn, Thr or Lys); Xaa at res.82 = (Ile, Val or Asn); Xaa at res.84 = (Lys or Arg); Xaa at res.85 = (Lys, Asn, Gln, His or Val); Xaa at res.86 = (Tyr or His); Xaa at res.87 = (Arg, Gln, Glu or Pro); Xaa at res.88 = (Asn,
25 Glu or Asp); Xaa at res.90 = (Val, Thr, Ala or Ile); Xaa at res.92 = (Arg, Lys, Val, Asp or Glu); Xaa at res.93 = (Ala, Gly, Glu or Ser); Xaa at res.95 = (Gly or Ala) and Xaa at res.97 = (His or Arg).

- 28 -

Generic Sequence 6

```

Cys Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Phe
  1                      5                      10
5  Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa
    15
    Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala
    20                      25
    Xaa Tyr Cys Xaa Gly Xaa Cys Xaa
10                      30                      35
    Xaa Pro Xaa Xaa Xaa Xaa Xaa
    40
    Xaa Xaa Xaa Asn His Ala Xaa Xaa
    45                      50
15  Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
    55
    Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
    60                      65
    Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa
20                      70
    Xaa Xaa Xaa Leu Xaa Xaa Xaa
    75                      80
    Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
    85
25  Xaa Xaa Xaa Xaa Met Xaa Val Xaa
    90                      95
    Xaa Cys Xaa Cys Xaa
    100

```

30 wherein each Xaa is independently selected from a group of one or more specified amino acids as defined by the following: "Res." means "residue" and Xaa at res.2 = (Lys, Arg, Ala or Gln); Xaa at res.3 = (Lys, Arg or

- 29 -

Met); Xaa at res.4 = (His, Arg or Gln); Xaa at res.5 =
(Glu, Ser, His, Gly, Arg, Pro, Thr, or Tyr); Xaa at
res.7 = (Tyr or Lys); Xaa at res.8 = (Val or Ile); Xaa
at res.9 = (Ser, Asp or Glu); Xaa at res.11 = (Arg,
5 Gln, Ser, Lys or Ala); Xaa at res.12 = (Asp, Glu, or
Lys); Xaa at res.13 = (Leu, Val or Ile); Xaa at res.16
= (Gln, Leu, Asp, His, Asn or Ser); Xaa at res.17 =
(Asp, Arg, Asn or Glu); Xaa at res.19 = (Ile or Val);
Xaa at res.20 = (Ile or Val); Xaa at res.21 = (Ala or
10 Ser); Xaa at res.23 = (Glu, Gln, Leu, Lys, Pro or Arg);
Xaa at res.24 = (Gly or Ser); Xaa at res.25 = (Tyr or
Phe); Xaa at res.26 = (Ala, Ser, Asp, Met, His, Gln,
Leu, or Gly); Xaa at res.28 = (Tyr, Asn or Phe); Xaa at
res.31 = (Glu, His, Tyr, Asp, Gln or Ser); Xaa at
15 res.33 = Glu, Lys, Asp, Gln or Ala); Xaa at res.35 =
(Ala, Ser, Pro, Gln or Asn); Xaa at res.36 = (Phe, Leu
or Tyr); Xaa at res.38 = (Leu, Val or Met); Xaa at
res.39 = (Asn, Asp, Ala, Thr or Pro); Xaa at res.40 =
(Ser, Asp, Glu, Leu, Ala or Lys); Xaa at res.41 = (Tyr,
20 Cys, His, Ser or Ile); Xaa at res.42 = (Met, Phe, Gly
or Leu); Xaa at res.43 = (Asn, Ser or Lys); Xaa at
res.44 = (Ala, Ser, Gly or Pro); Xaa at res.45 = (Thr,
Leu or Ser); Xaa at res.49 = (Ile, Val or Thr); Xaa at
res.50 = (Val, Leu or Ile); Xaa at res.51 = (Gln or
25 Arg); Xaa at res.52 = (Thr, Ala or Ser); Xaa at res.53
= (Leu or Ile); Xaa at res.54 = (Val or Met); Xaa at
res.55 = (His, Asn or Arg); Xaa at res.56 = (Phe, Leu,
Asn, Ser, Ala or Val); Xaa at res.57 = (Ile, Met, Asn,
Ala, Val or Leu); Xaa at res.58 = (Asn, Lys, Ala, Glu,
30 Gly or Phe); Xaa at res.59 = (Pro, Ser or Val); Xaa at
res.60 = (Glu, Asp, Gly, Val or Lys); Xaa at res.61 =
(Thr, Ala, Val, Lys, Asp, Tyr, Ser, Ala, Pro or His);
Xaa at res.62 = (Val, Ala or Ile); Xaa at res.63 = (Pro
or Asp); Xaa at res.64 = (Lys, Leu or Glu); Xaa at

- 30 -

res.65 = (Pro or Ala); Xaa at res.68 = (Ala or Val);
 Xaa at res.70 = (Thr, Ala or Glu); Xaa at res.71 =
 (Gln, Lys, Arg or Glu); Xaa at res.72 = (Leu, Met or
 Val); Xaa at res.73 = (Asn, Ser, Asp or Gly); Xaa at
 5 res.74 = (Ala, Pro or Ser); Xaa at res.75 = (Ile, Thr,
 Val or Leu); Xaa at res.76 = (Ser, Ala or Pro); Xaa at
 res.77 = (Val, Met or Ile); Xaa at res.79 = (Tyr or
 Phe); Xaa at res.80 = (Phe, Tyr, Leu or His); Xaa at
 res.81 = (Asp, Asn or Leu); Xaa at res.82 = (Asp, Glu,
 10 Asn or Ser); Xaa at res.83 = (Ser, Gln, Asn, Tyr or
 Asp); Xaa at res.84 = (Ser, Asn, Asp, Glu or Lys); Xaa
 at res.85 = (Asn, Thr or Lys); Xaa at res.87 = (Ile,
 Val or Asn); Xaa at res.89 = (Lys or Arg); Xaa at
 res.90 = (Lys, Asn, Gln, His or Val); Xaa at res.91 =
 15 (Tyr or His); Xaa at res.92 = (Arg, Gln, Glu or Pro);
 Xaa at res.93 = (Asn, Glu or Asp); Xaa at res.95 =
 (Val, Thr, Ala or Ile); Xaa at res.97 = (Arg, Lys, Val,
 Asp or Glu); Xaa at res.98 = (Ala, Gly, Glu or Ser);
 Xaa at res.100 = (Gly or Ala); and Xaa at res.102 =
 20 (His or Arg).

Particularly useful sequences for use as morphogens
 in this invention include the C-terminal domains, e.g.,
 the C-terminal 96-102 amino acid residues of Vgl,
 25 Vgr-1, DPP, OP-1, OP-2, CBMP-2A, CBMP-2B, GDF-1 (see
 Table II, below, and Seq. ID Nos. 5-14), as well as
 proteins comprising the C-terminal domains of 60A,
 BMP3, BMP5 and BMP6 (see Seq. ID Nos. 24-28), all of
 which include at least the conserved six or seven
 30 cysteine skeleton. In addition, biosynthetic

- 31 -

constructs designed from the generic sequences, such as COP-1, 3-5, 7, 16, disclosed in U.S. Pat. No. 5,011,691, also are useful. Other sequences include the inhibins/activin proteins (see, for example, U.S. Pat. Nos. 4,968,590 and 5,011,691). Accordingly, other useful sequences are those sharing at least 70% amino acid sequence homology or "similarity", and preferably 80% homology or similarity with any of the sequences above. These are anticipated to include allelic, species variants and other sequence variants (e.g., including "muteins" or "mutant proteins"), whether naturally-occurring or biosynthetically produced, as well as novel members of this morphogenic family of proteins. As used herein, "amino acid sequence homology" is understood to mean amino acid sequence similarity, and homologous sequences share identical or similar amino acids, where similar amino acids are conserved amino acids as defined by Dayoff et al., Atlas of Protein Sequence and Structure; vol.5, Suppl.3, pp.345-362 (M.O. Dayoff, ed., Nat'l BioMed. Research Fdn., Washington D.C. 1978.) Thus, a candidate sequence sharing 70% amino acid homology with a reference sequence requires that, following alignment of the candidate sequence with the reference sequence, 70% of the amino acids in the candidate sequence are identical to the corresponding amino acid in the reference sequence, or constitute a conserved amino acid change thereto. "Amino acid sequence identity" is understood to require identical amino acids between two aligned sequences. Thus, a candidate sequence sharing

- 32 -

60% amino acid identity with a reference sequence requires that, following alignment of the candidate sequence with the reference sequence, 60% of the amino acids in the candidate sequence are identical to the
5 corresponding amino acid in the reference sequence.

As used herein, all homologies and identities calculated use OP-1 as the reference sequence. Also as used herein, sequences are aligned for homology and
10 identity calculations using the method of Needleman et al. (1970) J.Mol. Biol. 48:443-453 and identities calculated by the Align program (DNASTar, Inc.) In all cases, internal gaps and amino acid insertions in the candidate sequence as aligned are ignored when making
15 the homology/identity calculation.

The currently most preferred protein sequences useful as morphogens in this invention include those having greater than 60% identity, preferably greater
20 than 65% identity, with the amino acid sequence defining the conserved six cysteine skeleton of hOP1 (e.g., residues 43-139 of Seq. ID No. 5). These most preferred sequences include both allelic and species variants of the OP-1 and OP-2 proteins, including the
25 *Drosophila* 60A protein. Accordingly, in another preferred aspect of the invention, useful morphogens include active proteins comprising species of polypeptide chains having the generic amino acid sequence herein referred to as "OPX", which
30 accommodates the homologies between the various identified species of OP1 and OP2 (Seq. ID No. 29).

- 33 -

In still another preferred aspect of the invention, useful morphogens include dimeric proteins comprising amino acid sequences encoded by nucleic acids that hybridize to DNA or RNA sequences encoding the C-terminal sequences defining the conserved seven cysteine domain of OP1 or OP2, e.g., nucleotides 1036-1341 and nucleotides 1390-1695 of Seq. ID No. 16 and 20, respectively, under stringent hybridization conditions. As used herein, stringent hybridization conditions are defined as hybridization in 40% formamide, 5 X SSPE, 5 X Denhardt's Solution, and 0.1% SDS at 37°C overnight, and washing in 0.1 X SSPE, 0.1% SDS at 50°C.

The morphogens useful in the methods, composition and devices of this invention include proteins comprising any of the polypeptide chains described above, whether isolated from naturally-occurring sources, or produced by recombinant DNA or other synthetic techniques, and includes allelic and species variants of these proteins, naturally-occurring or biosynthetic mutants thereof, as well as various truncated and fusion constructs. Deletion or addition mutants also are envisioned to be active, including those which may alter the conserved C-terminal cysteine skeleton, provided that the alteration does not functionally disrupt the relationship of these cysteines in the folded structure. Accordingly, such active forms are considered the equivalent of the specifically described constructs disclosed herein.

- 34 -

The proteins may include forms having varying glycosylation patterns, varying N-termini, a family of related proteins having regions of amino acid sequence homology, and active truncated or mutated forms of
5 native or biosynthetic proteins, produced by expression of recombinant DNA in host cells.

The morphogenic proteins can be expressed from intact or truncated cDNA or from synthetic DNAs in
10 procaryotic or eucaryotic host cells, and purified, cleaved, refolded, and dimerized to form morphogenically active compositions. Currently preferred host cells include E. coli or mammalian cells, such as CHO, COS or BSC cells. A detailed
15 description of the morphogens useful in the methods, compositions and devices of this invention is disclosed in international application US92/01968 (WO 92/15323) the disclosure of which is incorporated herein by reference.

20

Thus, in view of this disclosure, skilled genetic engineers can isolate genes from cDNA or genomic libraries of various different species which encode appropriate amino acid sequences, or construct DNAs
25 from oligonucleotides, and then can express them in various types of host cells, including both procaryotes and eucaryotes, to produce large quantities of active proteins useful as dietary compositions for enhancing tissue morphogenesis, including enhancing tissue
30 development and tissue viability in a variety of mammals, including humans.

- 35 -

The foregoing and other objects, features and advantages of the present invention will be made more apparent from the following detailed description of the invention.

- 36 -

Brief Description of the Drawings

5 The foregoing and other objects and features of this invention, as well as the invention itself, may be more fully understood from the following description, when read together with the accompanying drawings, in which:

10 FIG. 1A and B shows relative amounts of protein present in mammary gland extract eluate fractions of a C-18 reverse phase chromatography column (A), and the corresponding results of a Western Blot (B);

15 FIG. 2A and B shows relative amounts of protein present in bovine colostrum eluate fractions from purification scheme A of a C-18 reverse phase chromatography column (A), and the corresponding
20 results of a Western blot under reduced (1) and oxidized (2) conditions (B);

FIG. 3A and B shows relative amounts of protein present in bovine colostrum eluate fractions from
25 purification scheme B of a C-18 reverse phase chromatography column (A), and the corresponding results of a Western Blot under reduced conditions (B);

FIG. 4A and B shows relative amounts of protein
30 present in bovine 57 day milk eluate fractions of a C-18 reverse phase chromatography column (A), and the corresponding results of a Western Blot under reduced (1) and oxidized (2) conditions (B);

- 37 -

FIG. 5 shows Western Blot analysis of bovine colostrum using OP-1 and BMP2-specific antibodies;

FIG. 6A and B show results of in vivo and in vitro activity assays, respectively, for the corresponding fractions shown in Fig. 1;

FIG. 7 is a photomicrograph of an immunoblot showing the presence of hOP-1 in serum; and

FIG. 8A is a dose response curve for the induction of the 180 kDa and 140 kDa N-CAM isoforms in morphogen-treated NG108-15 cells;

FIG. 8B is a photomicrograph of a Western blot of whole cell extracts from morphogen-treated NG108-15 cells with an N-CAM-specific antibody; and

FIG. 9 (A and B) are photomicrographs showing the effect of morphogen-specific antibody on mouse development (9B) compared to untreated, control mice (9A).

- 38 -

Detailed Description of the Invention

It now has been discovered that the proteins
5 described herein are found in nursing mother's milk and
are useful as components of a dietary composition for
enhancing tissue morphogenesis in a mammal,
particularly in an individual at risk for normal tissue
development and viability. As described herein, these
10 proteins ("morphogens") are capable of enhancing tissue
development in growing mammals, stimulating CAM
expression and maintaining the normal tissue function
in adult tissue.

15 Provided below are detailed descriptions of
suitable morphogens useful in the compositions and
methods of this invention, as well as methods for their
administration and application, and numerous,
nonlimiting examples which demonstrate the suitability
20 of the morphogens described herein as active components
of a dietary composition for a mammal; and 2) provide
assays with which to test candidate morphogens for
their efficacy. Specifically, examples are provided
which (1) demonstrate the presence of endogenous
25 morphogen in milk and human serum (Examples 1 and 2),
(2) demonstrate the ability of morphogens to induce CAM
expression in a mammal (Example 3), (3) demonstrate the
ability of morphogens to enhance tissue development in
developing embryos (Example 4) and juveniles
30 (Example 5); (4) demonstrate the ability of morphogens
to reduce an osteoporotic condition in a mammal
(Example 6); (5) demonstrate the presence of morphogens
in developing tissues and adult stomach and gut tissue,
demonstrate the ability of parenterally provided

- 39 -

morphogen to localize to stomach tissue, and describe protocols for identifying morphogen-synthesizing tissue (Example 7) and (6) describe protocols for obtaining morphogen-specific antibodies and measuring morphogens in solution (Example 8).

I. Useful Morphogens

As defined herein a protein is morphogenic if it is capable of inducing the developmental cascade of cellular and molecular events that culminate in the formation of new, organ-specific tissue and comprises at least the conserved C-terminal six cysteine skeleton or its functional equivalent (see supra). Specifically, the morphogens generally are capable of all of the following biological functions in a morphogenically permissive environment: stimulating proliferation of progenitor cells; stimulating the differentiation of progenitor cells; stimulating the proliferation of differentiated cells; and supporting the growth and maintenance of differentiated cells. Details of how the morphogens useful in the method of this invention first were identified, as well as a description on how to make, use and test them for morphogenic activity are disclosed in international application US92/01968 (WO 92/15323). As disclosed therein, the morphogens may be purified from naturally-sourced material or recombinantly produced from procaryotic or eucaryotic host cells, using the genetic sequences disclosed therein. Alternatively, novel morphogenic sequences may be identified following the procedures disclosed therein.

- 40 -

Particularly useful proteins include those which comprise the naturally derived sequences disclosed in Table II. Other useful sequences include biosynthetic constructs such as those disclosed in U.S. Pat.

5 5,011,691, the disclosure of which is incorporated herein by reference (e.g., COP-1, COP-3, COP-4, COP-5, COP-7, and COP-16).

Accordingly, the morphogens useful in the methods
10 and compositions of this invention also may be described by morphogenically active proteins having amino acid sequences sharing 70% or, preferably, 80% homology (similarity) with any of the sequences described above, where "homology" is as defined herein
15 above.

The morphogens useful in the method of this invention also can be described by any of the 6 generic sequences described herein (Generic Sequences 1, 2, 3,
20 4, 5 and 6). Generic sequences 1 and 2 also may include, at their N-terminus, the sequence

Cys Xaa Xaa Xaa Xaa (Seq. ID No. 15)

1

5

25

Table II, set forth below, compares the amino acid sequences of the active regions of native proteins that have been identified as morphogens, including human OP-1 (hOP-1, Seq. ID Nos. 5 and 16-17), mouse OP-1
30 (mOP-1, Seq. ID Nos. 6 and 18-19), human and mouse OP-2 (Seq. ID Nos. 7, 8, and 20-23), CBMP2A (Seq. ID No. 9), CBMP2B (Seq. ID No. 10), BMP3 (Seq. ID No. 26), DPP (from Drosophila, Seq. ID No. 11), Vgl, (from Xenopus, Seq. ID No. 12), Vgr-1 (from mouse, Seq. ID No. 13),

- 41 -

GDF-1 (from mouse, Seq. ID Nos. 14, 32 and 33), 60A protein (from Drosophila, Seq. ID Nos. 24 and 25), BMP5 (Seq. ID No. 27) and BMP6 (Seq. ID No. 28). The sequences are aligned essentially following the method of Needleman et al. (1970) J. Mol. Biol., 48:443-453, calculated using the Align Program (DNASTar, Inc.) In the table, three dots indicates that the amino acid in that position is the same as the amino acid in hOP-1. Three dashes indicates that no amino acid is present in that position, and are included for purposes of illustrating homologies. For example, amino acid residue 60 of CBMP-2A and CBMP-2B is "missing". Of course, both these amino acid sequences in this region comprise Asn-Ser (residues 58, 59), with CBMP-2A then comprising Lys and Ile, whereas CBMP-2B comprises Ser and Ile.

TABLE II

20	hOP-1	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val
	mOP-1
	hOP-2	...	Arg	Arg
	mOP-2	...	Arg	Arg
25	DPP	...	Arg	Arg	...	Ser
	Vgl	Lys	Arg	His
	Vgr-1	Gly
	CBMP-2A	Arg	...	Pro
	CBMP-2B	...	Arg	Arg	...	Ser

- 42 -

	BMP3	...	Ala	Arg	Arg	Tyr	...	Lys	...	
	GDF-1	...	Arg	Ala	Arg	Arg	
	60A	...	Gln	Met	Glu	Thr	
	BMP5	
5	BMP6	...	Arg	
		1				5				
	hOP-1	Ser	Phe	Arg	Asp	Leu	Gly	Trp	Gln	Asp
10	mOP-1
	hOP-2	Gln	Leu	...
	mOP-2	Ser	Leu	...
	DPP	Asp	...	Ser	...	Val	Asp	...
	Vgl	Glu	...	Lys	...	Val	Asn
15	Vgr-1	Gln	...	Val
	CBMP-2A	Asp	...	Ser	...	Val	Asn	...
	CBMP-2B	Asp	...	Ser	...	Val	Asn	...
	BMP3	Asp	...	Ala	...	Ile	Ser	Glu
	GDF-1	Glu	Val	His	Arg
20	60A	Asp	...	Lys	His	...
	BMP5
	BMP6	Gln
			10					15		
25	hOP-1	Trp	Ile	Ile	Ala	Pro	Glu	Gly	Tyr	Ala
	mOP-1
	hOP-2	...	Val	Gln	Ser
	mOP-2	...	Val	Gln	Ser
	DPP	Val	Leu	Asp
30	Vgl	...	Val	Gln	Met
	Vgr-1	Lys
	CBMP-2A	Val	Pro	His

- 43 -

	CBMP-2B	Val	Pro	Gln
	BMP3	Ser	...	Lys	Ser	Phe	Asp
	GDF-1	...	Val	Arg	...	Phe	Leu
	60A	Gly
5	BMP5
	BMP6	Lys
				20					25	
10	hOP-1	Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ala
	mOP-1
	hOP-2	Ser
	mOP-2
	DPP	His	...	Lys	...	Pro
15	Vgl	...	Asn	Tyr	Pro
	Vgr-1	...	Asn	Asp	Ser
	CBMP-2A	...	Phe	His	...	Glu	...	Pro
	CBMP-2B	...	Phe	His	...	Asp	...	Pro
	BMP3	Ser	...	Ala	...	Gln
20	GDF-1	...	Asn	Gln	...	Gln
	60A	...	Phe	Ser	Asn
	BMP5	...	Phe	Asp	Ser
	BMP6	...	Asn	Asp	Ser
					30					35
25	hOP-1	Phe	Pro	Leu	Asn	Ser	Tyr	Met	Asn	Ala
	mOP-1
	hOP-2	Asp	...	Cys
	mOP-2	Asp	...	Cys
30	DPP	Ala	Asp	His	Phe	...	Ser
	Vgl	Tyr	Thr	Glu	Ile	Leu	...	Gly
	Vgr-1	Ala	His
	CBMP-2A	Ala	Asp	His	Leu	...	Ser
	CBMP-2B	Ala	Asp	His	Leu	...	Ser

- 44 -

5	GDF-1	Leu	...	Val	Ala	Leu	Ser	Gly	Ser**	...
	BMP3	Met	Pro	Lys	Ser	Leu	Lys	Pro
	60A	Ala	His
	BMP5	Ala	His	Met
	BMP6	Ala	His	Met
40										
10	hOP-1	Thr	Asn	His	Ala	Ile	Val	Gln	Thr	Leu
	mOP-1
	hOP-2	Leu	...	Ser	...
	mOP-2	Leu	...	Ser	...
	DPP	Val
15	Vgl	Ser	Leu
	Vgr-1
	CBMP-2A
	CBMP-2B
	BMP3	Ser	Thr	Ile	...	Ser	Ile
20	GDF-1	Leu	Val	Leu	Arg	Ala	...
	60A
	BMP5
	BMP6
		45					50			
25	hOP-1	Val	His	Phe	Ile	Asn	Pro	Glu	Thr	Val
	mOP-1	Asp
	hOP-2	...	His	Leu	Met	Lys	...	Asn	Ala	...
	mOP-2	...	His	Leu	Met	Lys	...	Asp	Val	...
	DPP	...	Asn	Asn	Asn	Gly	Lys	...
30	Vgl	Ser	...	Glu	Asp	Ile
	Vgr-1	Val	Met	Tyr	...
	CBMP-2A	...	Asn	Ser	Val	...	Ser	---	Lys	Ile
	CBMP-2B	...	Asn	Ser	Val	...	Ser	---	Ser	Ile
	BMP3	...	Arg	Ala**	Gly	Val	Val	Pro	Gly	Ile

- 45 -

	GDF-1	Met	...	Ala	Ala	Ala	...	Gly	Ala	Ala
	60A	Leu	Leu	Glu	...	Lys	Lys	...
	BMP5	Leu	Met	Phe	...	Asp	His	...
	BMP6	Leu	Met	Tyr	...
5			55					60		
	hOP-1	Pro	Lys	Pro	Cys	Cys	Ala	Pro	Thr	Gln
	mOP-1
10	hOP-2	Ala	Lys
	mOP-2	Ala	Lys
	DPP	Ala	Val
	Vgl	...	Leu	Val	Lys
	Vgr-1	Lys
15	CBMP-2A	Ala	Val	Glu
	CBMP-2B	Ala	Val	Glu
	BMP3	...	Glu	Val	...	Glu	Lys
	GDF-1	Asp	Leu	Val	...	Ala	Arg
	60A	Arg
20	BMP5	Lys
	BMP6	Lys
				65					70	
	hOP-1	Leu	Asn	Ala	Ile	Ser	Val	Leu	Tyr	Phe
25	mOP-1
	hOP-2	...	Ser	...	Thr	Tyr
	mOP-2	...	Ser	...	Thr	Tyr
	Vgl	Met	Ser	Pro	Met	...	Phe	Tyr
	Vgr-1	Val
30	DPP	...	Asp	Ser	Val	Ala	Met	Leu
	CBMP-2A	...	Ser	Met	Leu
	CBMP-2B	...	Ser	Met	Leu
	BMP3	Met	Ser	Ser	Leu	...	Ile	...	Phe	Tyr
	GDF-1	...	Ser	Pro	Phe	...

- 46 -

	60A	...	Gly	...	Leu	Pro	His
	BMP5
	BMP6
					75					80
5										
	hOP-1	Asp	Asp	Ser	Ser	Asn	Val	Ile	Leu	Lys
	mOP-1
	hOP-2	...	Ser	...	Asn	Arg
	mOP-2	...	Ser	...	Asn	Arg
10	DPP	Asn	...	Gln	...	Thr	...	Val
	Vgl	...	Asn	Asn	Asp	Val	...	Arg
	Vgr-1	Asn
	CBMP-2A	...	Glu	Asn	Glu	Lys	...	Val
	CBMP-2B	...	Glu	Tyr	Asp	Lys	...	Val
15	BMP3	...	Glu	Asn	Lys	Val
	GDF-1	...	Asn	...	Asp	Val	...	Arg
	60A	Leu	Asn	Asp	Glu	Asn
	BMP5
	BMP6	Asn
20						85				
	hOP-1	Lys	Tyr	Arg	Asn	Met	Val	Val	Arg	
	mOP-1	
25	hOP-2	...	His	Lys	
	mOP-2	...	His	Lys	
	DPP	Asn	...	Gln	Glu	...	Thr	...	Val	
	Vgl	His	...	Glu	Ala	...	Asp	
	Vgr-1	
30	CBMP-2A	Asn	...	Gln	Asp	Glu	
	CBMP-2B	Asn	...	Gln	Glu	Glu	

- 47 -

	BMP3	Val	...	Pro	Thr	...	Glu
	GDF-1	Gln	...	Glu	Asp	Asp
	60A	Ile	...	Lys
	BMP5
5	BMP6	Trp
		90					95		
	hOP-1	Ala	Cys	Gly	Cys	His			
10	mOP-1			
	hOP-2			
	mOP-2			
	DPP	Gly	Arg			
	Vgl	Glu	Arg			
15	Vgr-1			
	CBMP-2A	Gly	Arg			
	CBMP-2B	Gly	Arg			
	BMP3	Ser	...	Ala	...	Arg			
	GDF-1	Glu	Arg			
20	60A	Ser			
	BMP5	Ser			
	BMP6			
									100

25 **Between residues 56 and 57 of BMP3 is a Val residue;
 between residues 43 and 44 of GDF-1 lies
 the amino acid sequence Gly-Gly-Pro-Pro.

30 As is apparent from the foregoing amino acid
 sequence comparisons, significant amino acid changes
 can be made within the generic sequences while
 retaining the morphogenic activity. For example, while

- 48 -

the GDF-1 protein sequence depicted in Table II shares only about 50% amino acid identity with the hOP1 sequence described therein, the GDF-1 sequence shares greater than 70% amino acid sequence homology (or
5 "similarity") with the hOP1 sequence, where "homology" or "similarity" includes allowed conservative amino acid changes within the sequence as defined by Dayoff, et al., Atlas of Protein Sequence and Structure vol.5, supp.3, pp.345-362, (M.O. Dayoff, ed., Nat'l BioMed.
10 Res. Fd'n, Washington D.C. 1979.)

The currently most preferred protein sequences useful as morphogens in this invention include those having greater than 60% identity, preferably greater
15 than 65% identity, with the amino acid sequence defining the conserved six cysteine skeleton of hOP1 (e.g., residues 43-139 of Seq. ID No. 5). These most preferred sequences include both allelic and species variants of the OP-1 and OP-2 proteins, including the
20 *Drosophila* 60A protein. Accordingly, in still another preferred aspect, the invention includes morphogens comprising species of polypeptide chains having the generic amino acid sequence referred to herein as "OPX", which defines the seven cysteine skeleton and
25 accommodates the identities between the various identified mouse and human OP1 and OP2 proteins. OPX is presented in Seq. ID No. 29. As described therein, each Xaa at a given position independently is selected from the residues occurring at the corresponding
30 position in the C-terminal sequence of mouse or human OP1 or OP2 (see Seq. ID Nos. 5-8 and/or Seq. ID Nos. 16-23).

- 49 -

II. Formulations and Methods for Administering Therapeutic Agents

A. General Considerations

5

The morphogens may be provided to an individual by any suitable means, most preferably orally, or, alternatively, parenterally. Where the morphogen is to be provided parenterally, such as intravenously or by enteral feeding tube, the morphogen preferably comprises part of an aqueous solution. The solution is physiologically acceptable so that in addition to delivery of the desired morphogen to the patient, the solution does not otherwise adversely affect the patient's electrolyte and volume balance. The aqueous medium for the morphogen thus may comprise normal physiologic saline (0.85% NaCl, 0.15M), pH 7-7.4. The aqueous solution containing the morphogen can be made, for example, by dissolving the protein in 50% ethanol containing acetonitrile in 0.1% trifluoroacetic acid (TFA) or 0.1% HCl, or equivalent solvents. One volume of the resultant solution then is added, for example, to ten volumes of phosphate buffered saline (PBS), which further may include 0.1-0.2% human serum albumin (HSA). The resultant solution preferably is vortexed extensively. If desired, a given morphogen may be made more soluble by association with a suitable molecule. For example, the pro form of the morphogenic protein comprises a species that is soluble in physiologically buffered solutions. In fact, the endogenous protein is thought to be transported (e.g., secreted and circulated) in this form. This soluble form of the protein may be obtained from the culture medium of morphogen-secreting mammalian cells. Alternatively, a

30

- 50 -

soluble species may be formulated by complexing (e.g., via non-covalent interaction) the mature dimer (or an active fragment thereof) with part or all of one or, preferably, two pro domain peptides (see Section A.1, below). Another molecule capable of enhancing solubility and particularly useful for oral administrations, is casein, including derivatives and analogs thereof. For example, addition of 0.2% casein increases solubility of the mature active form of OP-1 in physiologically buffered solutions by 80%. Other components found in milk and/or various serum proteins also may be useful.

Useful solutions for parenteral administration may be prepared by any of the methods well known in the pharmaceutical art, described, for example, in Remington's Pharmaceutical Sciences (Gennaro, A., ed.), Mack Pub., 1990. Formulations may include, for example, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes, and the like. Biocompatible, preferably bioresorbable, polymers, including, for example, hyaluronic acid, collagen, polybutyrate, tricalcium phosphate, lactide and lactide/glycolide copolymers, may be useful excipients to control the release of the morphogen in vivo.

As described above, the dietary supplements comprising the morphogens described herein preferably are provided orally. Oral administration of proteins as therapeutics generally is not practiced as most proteins are readily degraded by digestive enzymes and acids in the mammalian digestive system before they can be absorbed into the bloodstream. However, the

- 51 -

morphogens described herein typically are acid stable and protease-resistant (see, for example, U.S. Pat.No. 4,968,590.) In addition, at least one morphogen, OP-1, has been identified in mammary gland extract, colostrum and 57-day milk (see Example 1, below). Moreover, the OP-1 purified from mammary gland extract is morphogenically active. Specifically, this protein induces endochondral bone formation in mammals when implanted subcutaneously in association with a suitable matrix material, using a standard in vivo bone assay, such as is disclosed in U.S. Pat.No. 4,968,590. Moreover, the morphogen also is detected in the bloodstream (see Example 2, below). These findings indicate that oral and parenteral administration are viable means for administering morphogens to an individual. In addition, while the mature forms of certain morphogens described herein typically are sparingly soluble, the morphogen form found in milk (and mammary gland extract and colostrum) is readily soluble, probably by association of the mature, morphogenically active form with part or all of at least one pro domain peptide and/or by association with one or more milk components. Accordingly, the compounds provided herein also may be associated with molecules capable of enhancing their solubility in vitro or in vivo.

The dietary compositions for oral administration may be formulated as a liquid, for example, as part of an aqueous medium as described above for parenteral administration, and which further may contain flavoring and coloring agents. The formulation also may be combined with a beverage or may be provided in a syrup. The dietary composition also may be provided as

- 52 -

an aerosol for oral or nasal administration.

Formulations for inhalation administration contain as excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-

5 lauryl ether, glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally.

Alternatively, the dietary composition may be provided as a solid, for example as a tablet, capsule or

10 lozenge. As for parenteral administration, formulations for oral administration also may include molecules to enhance a controlled release of the morphogen in vivo.

15 As will be appreciated by those skilled in the art, the concentration of the compounds described in a given dietary supplement composition will vary depending upon a number of factors, including the dosage number to be administered, the chemical characteristics (e.g.,
20 hydrophobicity) of the compounds employed, and the route of administration. The preferred dosage to be administered also is likely to depend on such variables as the type and extent of tissue development
25 enhancement desired, the type and extent of any tissue damage present to be repaired, the overall health status of the particular individual, the relative biological efficacy of the compound selected, the formulation of the compound excipients, and its route
30 of administration. In general terms, the compounds of this invention may be provided in a formulation containing about 0.001 to 10% w/v of morphogen to formulation. Typical dose ranges are from about 10 ng/kg to about 1 g/kg of body weight per day; a preferred dose range is from about 0.1 µg/kg to

- 53 -

100 mg/kg of body weight per day. Optimally, the morphogen dosage given is between 0.1-100 μ g of protein per kilogram weight of the individual. No obvious morphogen induced pathological lesions are induced when
5 mature morphogen (e.g., OP-1, 20 μ g) is administered daily to normal growing rats for 21 consecutive days. Moreover, 10 μ g systemic injections of morphogen (e.g., OP-1) injected daily for 10 days into normal newborn mice does not produce any gross abnormalities.

10

In administering morphogens parenterally in the methods of the present invention, preferably a large volume loading dose is used at the start of the treatment. The treatment then is continued with a
15 maintenance dose. In all cases administration dosages then can be monitored by measuring at intervals the levels of the morphogen in the blood.

A.1 Soluble Morphogen Complexes

20

A currently preferred form of the morphogen useful in therapeutic formulations, having improved solubility in aqueous solutions and consisting essentially of amino acids, is a dimeric morphogenic protein
25 comprising at least the 100 amino acid peptide sequence having the pattern of seven or more cysteine residues characteristic of the morphogen family complexed with a peptide comprising part or all of a pro region of a member of the morphogen family, or an allelic, species
30 or other sequence variant thereof. Preferably, the dimeric morphogenic protein is complexed with two peptides. Also, the dimeric morphogenic protein preferably is noncovalently complexed with the pro region peptide or peptides. The pro region peptides

- 54 -

also preferably comprise at least the N-terminal
eighteen amino acids that define a given morphogen
pro region. In a most preferred embodiment, peptides
defining substantially the full length pro region are
5 used.

Other soluble forms of morphogens include dimers of
the uncleaved pro forms of these proteins, as well as
"hemi-dimers" wherein one subunit of the dimer is an
10 uncleaved pro form of the protein, and the other
subunit comprises the mature form of the protein,
including truncated forms thereof, preferably
noncovalently associated with a cleaved pro domain
peptide.

15

As described above, useful pro domains include the
full length pro regions, as well as various truncated
forms hereof, particularly truncated forms cleaved at
proteolytic Arg-Xaa-Xaa-Arg cleavage sites. For
20 example, in OP-1, possible pro sequences include
sequences defined by residues 30-292 (full length
form); 48-292; and 158-292. Soluble OP-1 complex
stability is enhanced when the pro region comprises the
full length form rather than a truncated form, such as
25 the 48-292 truncated form, in that residues 30-47 show
sequence homology to the N-terminal portions of other
morphogens, and are believed to have particular utility
in enhancing complex stability for all morphogens.
Accordingly, currently preferred pro sequences are
30 those encoding the full length form of the pro region
for a given morphogen. Other pro sequences

- 55 -

contemplated to have utility include biosynthetic pro sequences, particularly those that incorporate a sequence derived from the N-terminal portion of one or more morphogen pro sequences.

5

As will be appreciated by those having ordinary skill in the art, useful sequences encoding the pro region may be obtained from genetic sequences encoding known morphogens. Alternatively, chimeric pro regions
10 can be constructed from the sequences of one or more known morphogens. Still another option is to create a synthetic sequence variant of one or more known pro region sequences.

15 In another preferred aspect, useful pro region peptides include polypeptide chains comprising an amino acid sequence encoded by a nucleic acid that hybridizes under stringent conditions with a DNA or RNA sequence encoding at least the N-terminal eighteen amino acids
20 of the pro region sequence for OP1 or OP2, e.g., nucleotides 136-192 and 152-211 of Seq. ID No. 16 and 20, respectively.

25 A.1A Isolation of Soluble morphogen complex from conditioned media or body fluid

Morphogens are expressed from mammalian cells as soluble complexes. Typically, however the complex is disassociated during purification, generally by
30 exposure to denaturants often added to the purification solutions, such as detergents, alcohols, organic solvents, chaotropic agents and compounds added to reduce the pH of the solution. Provided below is a currently preferred protocol for purifying the soluble

- 56 -

proteins from conditioned media (or, optionally, a body fluid such as serum, cerebro-spinal or peritoneal fluid), under non-denaturing conditions. The method is rapid, reproducible and yields isolated soluble morphogen complexes in substantially pure form.

Soluble morphogen complexes can be isolated from conditioned media using a simple, three step chromatographic protocol performed in the absence of denaturants. The protocol involves running the media (or body fluid) over an affinity column, followed by ion exchange and gel filtration chromatographies. The affinity column described below is a Zn-IMAC column. The present protocol has general applicability to the purification of a variety of morphogens, all of which are anticipated to be isolatable using only minor modifications of the protocol described below. An alternative protocol also envisioned to have utility an immunoaffinity column, created using standard procedures and, for example, using antibody specific for a given morphogen pro domain (complexed, for example, to a protein A-conjugated Sepharose column.) Protocols for developing immunoaffinity columns are well described in the art, (see, for example, Guide to Protein Purification, M. Deutscher, ed., Academic Press, San Diego, 1990, particularly sections VII and XI.)

In this experiment OP-1 was expressed in mammalian CHO (chinese hamster ovary) cells as described in the art (see, for example, international application US90/05903 (WO91/05802).) The CHO cell conditioned media containing 0.5% FBS was initially purified using Immobilized Metal-Ion Affinity Chromatography (IMAC).

- 57 -

The soluble OP-1 complex from conditioned media binds very selectively to the Zn-IMAC resin and a high concentration of imidazole (50 mM imidazole, pH 8.0) is required for the effective elution of the bound

5 complex. The Zn-IMAC step separates the soluble OP-1 from the bulk of the contaminating serum proteins that elute in the flow through and 35 mM imidazole wash fractions. The Zn-IMAC purified soluble OP-1 is next applied to an S-Sepharose cation-exchange column

10 equilibrated in 20 mM NaPO_4 (pH 7.0) with 50 mM NaCl. This S-Sepharose step serves to further purify and concentrate the soluble OP-1 complex in preparation for the following gel filtration step. The protein was applied to a Sephacryl S-200HR column equilibrated in

15 TBS. Using substantially the same protocol, soluble morphogens also may be isolated from one or more body fluids, including serum, cerebro-spinal fluid or peritoneal fluid.

20 IMAC was performed using Chelating-Sepharose (Pharmacia) that had been charged with three column volumes of 0.2 M ZnSO_4 . The conditioned media was titrated to pH 7.0 and applied directly to the Zn-IMAC resin equilibrated in 20 mM HEPES (pH 7.0) with 500 mM

25 NaCl. The Zn-IMAC resin was loaded with 80 mL of starting conditioned media per mL of resin. After loading, the column was washed with equilibration buffer and most of the contaminating proteins were eluted with 35 mM imidazole (pH 7.0) in equilibration

30 buffer. The soluble OP-1 complex then is eluted with 50 mM imidazole (pH 8.0) in 20 mM HEPES and 500 mM NaCl.

- 58 -

The 50 mM imidazole eluate containing the soluble OP-1 complex was diluted with nine volumes of 20 mM NaPO_4 (pH 7.0) and applied to an S-Sepharose (Pharmacia) column equilibrated in 20 mM NaPO_4 (pH 7.0) with 50 mM NaCl. The S-Sepharose resin was loaded with an equivalent of 800 mL of starting conditioned media per mL of resin. After loading the S-Sepharose column was washed with equilibration buffer and eluted with 100 mM NaCl followed by 300 mM and 500 mM NaCl in 20 mM NaPO_4 (pH 7.0). The 300 mM NaCl pool was further purified using gel filtration chromatography. Fifty mls of the 300 mM NaCl eluate was applied to a 5.0 X 90 cm Sephacryl S-200HR (Pharmacia) equilibrated in Tris buffered saline (TBS), 50 mM Tris, 150 mM NaCl (pH 7.4). The column was eluted at a flow rate of 5 mL/minute collecting 10 mL fractions. The apparent molecular of the soluble OP-1 was determined by comparison to protein molecular weight standards (alcohol dehydrogenase (ADH, 150 kDa), bovine serum albumin (BSA, 68 kDa), carbonic anhydrase (CA, 30 kDa) and cytochrome C (cyt C, 12.5 kDa). The purity of the S-200 column fractions was determined by separation on standard 15% polyacrylamide SDS gels stained with coomassie blue. The identity of the mature OP-1 and the pro-domain was determined by N-terminal sequence analysis after separation of the mature OP-1 from the pro-domain using standard reverse phase C18 HPLC.

The soluble OP-1 complex elutes with an apparent molecular weight of 110 kDa. This agrees well with the predicted composition of the soluble OP-1 complex with one mature OP-1 dimer (35-36 kDa) associated with two

- 59 -

pro-domains (39 kDa each). Purity of the final complex can be verified by running the appropriate fraction in a reduced 15% polyacrylamide gel.

- 5 The complex components can be verified by running the complex-containing fraction from the S-200 or S-200HR columns over a reverse phase C18 HPLC column and eluting in an acetonitrile gradient (in 0.1% TFA), using standard procedures. The complex is dissociated
- 10 by this step, and the pro domain and mature species elute as separate species. These separate species then can be subjected to N-terminal sequencing using standard procedures (see, for example, Guide to Protein Purification, M. Deutscher, ed., Academic
- 15 Press, San Diego, 1990, particularly pp. 602-613), and the identity of the isolated 36kD, 39kDa proteins confirmed as mature morphogen and isolated, cleaved pro domain, respectively. N-terminal sequencing of the isolated pro domain from mammalian cell produced OP-1
- 20 revealed 2 forms of the pro region, the intact form (beginning at residue 30 of Seq. ID No. 16) and a truncated form, (beginning at residue 48 of Seq. ID No. 16.) N-terminal sequencing of the polypeptide subunit of the isolated mature species reveals a range of
- 25 N-termini for the mature sequence, beginning at residues 293, 300, 313, 315, 316, and 318, of Seq. ID No. 16, all of which are active as demonstrated by the standard bone induction assay.

- 60 -

A.1B. In Vitro Soluble Morphogen Complex Formation

As an alternative to purifying soluble complexes from culture media or a body fluid, soluble complexes may be formulated from purified pro domains and mature dimeric species. Successful complex formation apparently requires association of the components under denaturing conditions sufficient to relax the folded structure of these molecules, without affecting disulfide bonds. Preferably, the denaturing conditions mimic the environment of an intracellular vesicle sufficiently such that the cleaved pro domain has an opportunity to associate with the mature dimeric species under relaxed folding conditions. The concentration of denaturant in the solution then is decreased in a controlled, preferably step-wise manner, so as to allow proper refolding of the dimer and pro regions while maintaining the association of the pro domain with the dimer. Useful denaturants include 4-6M urea or guanidine hydrochloride (GuHCl), in buffered solutions of pH 4-10, preferably pH 6-8. The soluble complex then is formed by controlled dialysis or dilution into a solution having a final denaturant concentration of less than 0.1-2M urea or GuHCl, preferably 1-2 M urea or GuHCl, which then preferably can be diluted into a physiological buffer. Protein purification/renaturing procedures and considerations are well described in the art, and details for developing a suitable renaturing protocol readily can be determined by one having ordinary skill in the art.

- 61 -

One useful text on the subject is Guide to Protein Purification, M. Deutscher, ed., Academic Press, San Diego, 1990, particularly section V. Complex formation also may be aided by addition of one or more chaperone proteins.

A.1C. Stability of Soluble Morphogen Complexes

The stability of the highly purified soluble morphogen complex in a physiological buffer, e.g., tris-buffered saline (TBS) and phosphate-buffered saline (PBS), can be enhanced by any of a number of means. Currently preferred is by means of a pro region that comprises at least the first 18 amino acids of the pro sequence (e.g., residues 30-47 of Seq. ID NO. 16 for OP-1), and preferably is the full length pro region. Residues 30-47 show sequence homology to the N-terminal portion of other morphogens and are believed to have particular utility in enhancing complex stability for all morphogens. Other useful means for enhancing the stability of soluble morphogen complexes include three classes of additives. These additives include basic amino acids (e.g., L-arginine, lysine and betaine); nonionic detergents (e.g., Tween 80 or Nonidet P-120); and carrier proteins (e.g., serum albumin and casein). Useful concentrations of these additives include 1-100 mM, preferably 10-70 mM, including 50 mM, basic amino acid; 0.01-1.0%, preferably 0.05-0.2%, including 0.1% (v/v) nonionic detergent; and 0.01-1.0%, preferably 0.05-0.2%, including 0.1% (w/v) carrier protein.

- 62 -

B. Considerations for Infant and Other Formulas

1. Infant Formulas

5 In all cases the morphogens of this invention preferably are added to an infant formula that complies with the nutritional guidelines provided by the AAP and ESPGAN. Basic ingredients for infant formulas include cow's milk, protein, whey proteins, casein and its
10 salts (i.e. calcium caseinate). Soy protein isolates may be substituted for milk-derived proteins, and preferably are used in the products made for infants with lactose intolerance and/or cow's protein intolerance. Protein hydrolyzates (i.e. casein and
15 lactalbumin hydrolyzates) with low molecular weight, also may be used for these products.

 The proportions of the diverse component nutrients preferably are similar to those of human milk. Thus,
20 the ratio of whey proteins to casein preferably varies from 60:40 to 70:30 in infant formulas based on milk. The mixture of fats employed is made up of edible fats to provide an essential fatty acid profile. Lactose preferably is used as the carbohydrate source for
25 at-term newborns infants, and dextrinmaltose preferably is employed in products used for the treatment of lactose intolerance and malabsorption syndromes in infancy.

30 Infant formulas according to the invention also preferably contain minerals (including calcium, phosphorus, sodium, potassium, chloride, magnesium, iron, zinc, copper, manganese and iodine) and vitamins (including vitamin A, vitamin D3, vitamin C, vitamin

- 63 -

B1, vitamin B2, vitamin B6, vitamin B12, pantothenic acid, vitamin E, vitamin K1, folic acid, biotin) adequate for the infants' requirements. Also, in the products whose source of proteins is derived from soy
5 or protein isolates or hydrolyzates, carnitine preferably is included to satisfy the nutritional requirements for this compound in infants with malabsorptive syndromes.

10 A typical ready-to-feed morphogen-enriched formulation for infants, when diluted to feeding concentrations, preferably comprises in addition to the added morphogen, from about 1-5% by weight fat, from about 0.01 to about 0.5% by weight immunoglobulins as
15 appropriate, from about 4-10% by weight carbohydrate in a quantity substantially to mimic the carbohydrate content of human mother's milk, from about 0.5 to 4% by weight protein in a quantity substantially to mimic the protein content of human mother's milk, optional
20 vitamins and minerals as required, a total solids content of from about 8 to 17% by weight, and the remainder water.

A typical protein source for use in infant formula
25 is electrodialyzed whey or electrodialyzed skim milk or milk whey, although other protein sources are also available and may be used. Preferred sugars include food grade substances such as glucose, dextrose, sucrose, or edible lactose. The following vitamins and
30 minerals may also be incorporated in the infant formula: calcium, phosphorus, potassium, sodium, chloride, magnesium, manganese, iron, copper, zinc, selenium, iodine, and vitamins A, E, D, and B complex.

- 64 -

These micronutrients are added in the form of commonly accepted nutritional compounds in amounts equivalent to those present in human milk on a per calories basis.

5 The infant formula according to the present invention also preferably is sterilized and subsequently used on a ready-to-feed basis, or can be stored as a concentrate. The concentrate can be prepared using standard procedures known in the art,
10 and the formula can be reconstituted by rehydrating the concentrate. The infant formula preferably is a stable liquid and has a suitable shelf life. A more detailed description of infant formula considerations, including preferred formulations for newborn, preterm and low
15 birth-weight infants, lactose-intolerant infants, may be found, for example, in US Pat. No. 5,066,500 to Gil et al., the disclosure of which is incorporated herein by reference.

20 2. Other Nutritional Products

 The morphogen-enriched dietary products for balanced nutrition (e.g., dietary food formulations) according to the present invention, preferably have, in
25 addition to added morphogen, a composition of nutrients adequate to the specific requirements of not only healthy human in need of a balanced nutritional product, but also those individuals at risk for lost or reduced tissue function due malnutrition-maladsorption
30 disorder, and/or altered metabolism. Individuals particularly affected by an altered metabolic function include postmenopausal women or aged individuals, hypercatabolic individuals, and individuals undergoing periods of rapid growth or physical stress, such as

- 65 -

developing juveniles, and pregnant, lactating and nursing mothers. Other individuals at risk are those suffering from malnutrition, induced, for example, by starvation and/or an eating disorder, and individuals
5 affected with energy-protein malnutrition and in hypercatabolic states derived from traumatic, septic, surgical processes and other clinically-derived malabsorption syndromes.

- 10 Morphogen-enriched nutritional products according to the present invention preferably also provide mineral elements which include trace elements and vitamins in adequate proportions to satisfy the specific requirements of normal healthy individuals as
15 well as individuals at risk, such as those suffering malabsorption-malnutrition processes and in a hypercatabolic state. The nutritional products also preferably are enriched with amino acids sources, vitamins, nucleosides and/or nucleotides in similar
20 amounts to those present in ordinary foods.

- As described above for infant formulas, liquid products may be formulated ready for consumption or as concentrates to be diluted before use. Preferably,
25 liquid dietary compositions have pH values generally ranging from about 6.0 to about 8.0, most preferably 6.8-7.5.

- 66 -

Useful dietary compositions and considerations for their formulation are well described in the medical and nutritional arts. Useful compositions for clinical nutrition, also are described in detail in US

5 Pat.No. 5,066,500.

III. Examples

10 Example 1. Determination of the Presence of Morphogen in Milk

Morphogenically active OP-1 was demonstrated to be present in mammary gland extract, colostrum, and milk, as described below. The discovery that the morphogen naturally is present in milk, together with the known observation that mature, active OP-1 is acid-stable and protease-resistant, indicate that oral administration is a useful route for therapeutic administration of morphogen to a mammal. Oral administration typically is the preferred mode of delivery for extended or prophylactic therapies. In addition, the identification of morphogen in all milk forms, including colostrum, indicates that the protein plays a significant role in tissue development, including skeletal development of juveniles.

15
20
25

Rat mammary gland extract and bovine colostrum and 57 day milk were subjected to purification procedures designed to partially purify OP-1. The partially purified product then was examined for the presence of OP-1 by Western blot analysis using OP-1-specific antisera, and tested for in vivo and in vitro activity.

30

- 67 -

1.1 Purification

The purification protocol for all three "milk" forms (e.g., mammary gland extract, colostrum and 57-day milk), involved three chromatography steps: (1) cation-exchange chromatography (S-Sepharose and followed by Phenyl-Sepharose chromatography); (2) Copper-Immobilized Metal Affinity chromatography (Cu⁺⁺-IMAC); and finally, (3) C-18 reverse phase chromatography. Fractions were sampled at each step for the presence of OP-1. Fraction samples for testing were dialyzed versus water/0.1% TFA, then against 30% acetonitrile/0.1% TFA for analysis on SDS-polyacrylamide gels and immunoblots, using standard methodologies well described in the art. Unless otherwise stated, the primary antibody used for the immunoblots was made against full length OP-1 produced in E.coli using standard recombinant DNA and antibody production techniques (see, for example, Example 8, below for a general description for producing morphogen-specific antibodies.) Fractions found to contain the morphogen then were applied to the next column step or used in the immunoreactivity or activity assays described below.

25

Essentially the same protocol was followed for all three milk sources, except that two alternative cation-exchange methodologies were employed for colostrum purification, described in detail below. Unless

- 68 -

otherwise indicated, all chemicals referenced are standard, commercially available reagents, readily available from a number of sources, including Sigma Chemical, Co., St. Louis; Calbiochem, Corp., San Diego
5 and Aldrich Chemical Co., Milwaukee.

step 1. Cation-Exchange Chromatography

The S-Sepharose purification step was performed as
10 follows. 200ml of cation exchanger (S-Sepharose, Sigma Chemical Corp.) were equilibrated with equilibration buffer (6M urea, 20mM MES, 70mM NaCl, pH 6.5). The supernatant from the centrifuged extract was diluted to final concentration of 6M urea, 20mM MES, 70mM NaCl, pH
15 6.5. After loading, the column was washed to baseline using equilibration buffer, and the bound components were eluted stepwise from the column with 6M urea, 20mM MES, 100mM and 500mM NaCl, pH 6.5. The more tightly bound components then were eluted with 4M guanidine,
20 20mM sodium phosphate, pH 7.0.

The Phenyl-Sepharose purification step was performed as follows. 15ml of Phenyl-Sepharose CL-4B (Sigma) were equilibrated with 6M urea, 20mM HEPES, 1M
25 ammonium sulfate, 300mM NaCl, pH 7.0. The 500mM NaCl eluate from the S-Sepharose step was diluted with 6M urea, 20mM HEPES, 3M ammonium sulfate, 300mM NaCl, pH 7.0, to a final concentration of 1M ammonium sulfate, pH 7.0. After loading, the column was washed to
30 baseline with equilibration buffer. The column was eluted with 6M urea, 20mM HEPES, 0.6M ammonium sulfate, 300mM NaCl, pH 7.0, and then with 4M guanidine, 20mM sodium phosphate, pH 7.0.

- 69 -

Two alternative cation-exchange chromatography schemes (A and B) were exploited in the purification of OP-1 from colostrum, as follows. For both schemes, 200 ml of S-Sepharose (Sigma) was poured into a 5 X 10 cm
5 Bio-Rad econocolumn (Bio-Rad, Inc. Cambridge.)

Scheme A: The colostrum, which had been diluted to 6M urea, 20mM sodium phosphate, pH 7.0, was loaded onto a column equilibrated with 6M urea, 20mM sodium
10 phosphate, 50mM NaCl, pH 7.0. Elution was stepwise, with 6M urea, 20mM sodium phosphate, 100mM and then 500mM NaCl, pH 7.0; and the final wash was with 4M guanidine, 20mM sodium phosphate, pH 7.0. The Phenyl-Sepharose column was run as described above, except
15 that sodium phosphate was used as the running buffer instead of HEPES. The Phenyl-Sepharose bound fraction (0.0M ammonium sulfate eluate) from scheme A then was dialyzed into 6M urea, 20mM Hepes, 500mM NaCl, pH 7.0, before it was applied to the Cu⁺⁺-IMAC column, which
20 was run as described below.

Scheme B: The alternative S-Sepharose purification was performed as follows. Ethanol-precipitated protein was loaded onto an S-Sepharose column equilibrated in
25 6M urea, 20mM MES, 50mM NaCl, pH 6.5. Elution was stepwise with 6M urea, 20mM MES, 100mM NaCl and then 500mM NaCl, and finally 4M guanidine, 20mM sodium phosphate, pH 7.0. The Phenyl-Sepharose column was run as described above, with the 0.0M ammonium sulfate
30 eluate then applied to a Cu⁺⁺-IMAC column.

- 70 -

step 2. Cu++IMAC Chromatography

The Cu++IMAC purification step was performed as follows. 10ml of Pharmacia Fast Flow Chelating Resin
5 were charged with 0.2M cupric sulfate, and equilibrated with 6M urea, 20mM HEPES, 0.5M NaCl, pH 7.0. After loading, the column was washed to baseline with equilibration buffer. Elution from the column was stepwise, using equilibration buffer containing 1mM,
10 5mM, or 10mM imidazole. The column then was stripped with equilibration buffer containing 10mM EDTA. The 10mM imidazole elution was dialyzed against water/0.1% TFA, then against 30% acetonitrile/0.1% TFA.

15 step 3. Reverse Phase Chromatography

The C-18 reverse phase chromatography purification step was performed as follows. A HPLC C-18 semi-prep column was used for the final purification step. The
20 gradient used was 30-50% acetonitrile/0.1% TFA over 60 minutes at 3ml/minutes. After the sample was loaded, the column was washed to baseline with 30% acetonitrile/0.1% TFA before the gradient is started. Fractions collected were 3ml in size. Chromatograms
25 were read at 214 nm.

(a) OP-1 from Rat Mammary Gland Extract

Mammary glands were obtained from 2 female Long
30 Evan rats (Charles River Labs, Wilmington, MA) one week post-partum. The excised glands were mildly homogenized in 6M urea, 20mM methylethansulfonate

- 71 -

(MES), 0.5M NaCl, pH 6.5 using a polytron homogenizer. The suspension then was centrifuged for 20 minutes at 8,000 RPM, and the supernatant removed for further purification.

5

Following S-Sepharose chromatography, fractions containing 6M urea, 20mM MES containing 500mM NaCl, also appeared to contain OP-1 as determined by SDS and immunoblot, and were applied to the Phenyl-Sepharose
10 column. The eluate from the 6M urea, 20mM HEPES, 300mM NaCl, pH 7.0 elution step from this column were found to contain OP-1. This eluate then was applied to a Cu⁺⁺-IMAC column. Eluate fractions found to contain OP-1 were then applied to the C-18 column and
15 chromatographed as described.

Figure 1(A) shows the chromatogram and 1(B) the corresponding Western blot for fractions from the C-18 reverse phase chromatography step run under reducing
20 conditions. Lane S of the Western blot is a standard, containing reduced, purified, recombinantly-produced OP-1. The arrows show molecular weight markers corresponding to 17, 27, and 39 Kd. The reduced monomer run at approximately 16-18 Kd; the oxidized homodimer
25 at approximately 36 Kd. Lanes 13-30 represent the corresponding fractions of the C-18 reverse phase column as numbered in Fig.1(A). As can be seen in Fig.1(B), mammary extract OP-1 elutes primarily in fractions 21-25 from this final chromatography step.

30

- 72 -

(b) OP-1 from Bovine Colostrum

Colostrum is the first milk to be produced by the mother following birth. Approximately 5 gallons of
5 bovine colostrum were obtained from a local dairy farm and delipidated by centrifugation (8000 rpm for approximately 10 min. at 4°C). The supernatant then was filtered through cheese cloth. The filtered supernatant was stored in 500ml aliquots at 70°C.

10

50 ml of colostrum were diluted with 100ml of 9M urea, 30mM sodium phosphate, pH 7.0. Alternately, 50ml of colostrum was added to 50ml of 8M guanidine-HCl, 50mM Tris, pH 7.2 and precipitated with 40%, then 85%
15 ice cold ethanol. The pellet was washed with 90% cold ethanol and lyophilized overnight. The lyophilized pellet was resuspended in 6M urea, 20mM MES, 500mM NaCl, pH 6.5, stirred overnight at 4°C, and centrifuged at 9,000 RPM for 10 minutes to clarify the suspension
20 before loading onto the column as described in schemes A and B, above.

Following S-Sepharose chromatography by scheme A, both the 100mM and the 500 mM eluate fractions were
25 found to contain OP-1, with the 100mM fraction containing relatively more morphogen. This fraction then was loaded onto the Phenyl-Sepharose column following dilution with an equal volume of 6M urea, 20 mM sodium phosphate, 2M ammonium sulfate, and 300mM
30 NaCl.

- 73 -

Following S-Sepharose chromatography by scheme B, the 500mM NaCl eluate was found to contain OP-1 and was loaded onto a Phenyl-Sepharose column as described above, following dilution with 6M urea, 40mM HEPES, 2M ammonium sulfate, pH 7.0.

Following Cu++IMAC chromatography OP-1 was identified in the 5mM and 10mM imidazole eluates for both purification schemes, and was dialyzed for further purification on the C-18 column.

Both purification schemes produce purified OP-1, as determined by immunoblot. Figure 2 shows the chromatogram (A) and corresponding Western blot (B) for results of purification scheme A (Fig. 2B-1, reduced and Fig. 2B-2, oxidized); and Figure 3 shows the chromatogram (A) and Western blot (B, reduced) for C-18-purified protein from scheme B. As for Fig. 1B, lane S in Figs. 2B and 3B is a standard, containing purified, recombinantly produced OP-1; 17, 27 and 39 are molecular weight markers, and lane numbers correspond to fraction numbers in the corresponding chromatograms. OP-1 purified by scheme A appears predominantly in fractions 18-27, and OP-1 purified by scheme B appears predominantly in fractions 18-25.

OP-1 from 57-day milk

Milk was obtained from the same cow from which the colostrum came, 57 days after the birth of the calf. The milk was delipidated by centrifugation at 10,000 RPM for 15 minutes, and the milk was poured off and away from the fat layer.

- 74 -

100ml of milk then were diluted with 200ml of 9M urea, 30mM MES, pH 6.5 and loaded onto a 200ml S-Sepharose column which had been equilibrated with 6M urea, 20mM MES, 50mM NaCl, pH 6.5. Elution was with 6M urea, 20mM MES, 100mM and 500mM NaCl, and 4M guanidine, 20mM sodium phosphate, pH 7.0. The 500mM elution was put over a Phenyl-Sepharose column after being diluted with an equal volume of 6M urea, 20mM MES, 2M ammonium sulfate, pH 7.0.

10

The Phenyl-Sepharose column then was run as described above. The Phenyl-Sepharose-bound sample was eluted and applied to a Cu++IMAC column, prepared and run as described above. The 10mM imidazole eluate was found to contain OP-1 and was dialyzed for further purification on the C-18 column.

The C-18 reverse phase chromatography column and gradient were performed as described above. The results are presented in Fig. 4A (chromatogram) and 4B (immunoblot, 10B-1, oxidized; 4B-2, reduced.) As above, lane S is a standard, containing purified, recombinantly produced OP-1; 17, 27, and 39 are molecular weight markers, and the lane numbers correspond to the fractions numbers in Fig. 4A. OP-1 purified from 57-day milk appears predominantly in fractions 18-26.

1.2 OP-1 Characterization by immunoreactivity

30

OP-1 purified from the different milk sources as described above also were characterized by Western blotting using antibodies raised against OP-1 and BMP2. Antibodies were prepared using standard immunology

- 75 -

protocols well known in the art, and as described in Example 8, below using full-length E. coli-produced OP-1 and BMP2 as the immunogens.

5 As shown in Fig. 5 OP-1 purified from colostrum reacts with the anti-OP-1 antibody, but not with anti-BMP2 antibody. In Fig. 5A and B, lane 1 contains reduced, purified, recombinantly-produced OP-1; lane 2 contains C-18 purified bovine colostrum, and lane 3
10 contains reduced COP-16, a biosynthetic construct having morphogenic activity and an amino acid sequence modeled on the proteins described herein, but having highest amino acid sequence homology with BMP2 (see US Pat. No. 5,011,691 for the COP-16 amino acid sequence.)
15 In Fig. 5A the gel was probed with anti-OP-1 antibody; in Fig. 5B, the gel was probed with anti-BMP2 antibody. As can be seen in the figure, anti-OP-1 antibody hybridizes with protein in lanes 1 and 2, but not 3; while anti-BMP2 antibody hybridizes with lane 3 only.

20

C-18 purified mammary gland extract and 57-day milk also were shown to react with anti-OP-1 antibodies, including antibody raised against the full length E. coli OP-1, full length mammalian-produced OP-1, and
25 the OP-1 Ser-17-Cys peptide (e.g., the OP-1 N-terminal 17 amino acids).

1.3 OP-1 Characterization by Activity

30 The morphogenic activity of OP-1 purified from mammary gland extract was evaluated in vivo as follows. 33% of each OP-1 immunoreactive fraction of C-18-purified mammary gland extract was lyophilized and resuspended in 220 μ l of 50% acetonitrile/0.1% TFA. After

- 76 -

vortexing, 25 mg of collagen matrix was added. The samples were lyophilized overnight, and implanted in Long Evans rats (Charles River Laboratories, Wilmington, MA, 28-35 days old). Each fraction was
5 implanted in duplicate. For details of the collagen matrix implantation procedure, see, for example, U.S. Pat. No. 4,968,590, hereby incorporated by reference. After 12 days, the implants were removed and evaluated for new bone formation by histological observation.

10

The results are presented in Fig. 6A, where "% activity" refers to the % of bone formation/total area covered by bone in the histology sample. In the figure, solid bars represent implants using mammary
15 extract-derived OP-1, where the fraction numbers correspond to the related fractions eluted from the C-18 reverse phase column (see Fig. 1B), and the hatched bar represents implants using recombinantly produced OP-1 (600 ng). The results demonstrate that
20 the peak bone forming activity of C-18-purified mammary gland extract corresponds with the immunoreactive fraction peaks of Fig. 1B (compare Fig. 6A and 1B.)

Similarly, the morphogenic activity of OP-1
25 purified from mammary gland extract was evaluated in vitro by measuring alkaline phosphatase activity in vitro using the following assay. Test samples were prepared using 15-20% of individual immunoreactive fractions from the C-18 run which were precipitated and
30 resuspended in a smaller volume of 50% acetonitrile/0.1% TFA. Alkaline phosphatase activity was tested using ROS 17/2.8 cells (Rat Osteosarcoma, e.g., obtained, for example, from Dr. Robert J. Majeska, Mt. Sinai Medical Center, New York, NY, in a

- 77 -

standard alkaline phosphatase activity assay (see, for example, U.S. Pat. No. 4,968,590). The results, presented in Fig. 6B, indicate that the immunoreactive fractions obtained from C-18-purified mammary gland extract correspond with alkaline phosphatase activity in vitro (compare Fig. 6B and Fig. 1B.) In Fig. 6B solid bars represent assays performed with mammary gland-purified OP-1, where the fraction numbers correspond to the related fractions eluted from the C-18 reverse phase column (see Fig. 1B), the hatched bar represents the assay performed with purified, recombinantly-produced OP-1 (100ng/ml), and the cross-hatched bar represents background. As for Fig. 6A, alkaline phosphatase activity corresponds with immunoreactivity of the C-18-purified extract (compare Fig. 6B and 1B.)

Example 2. Morphogen Identification in Human Serum

OP-1 was detected in human serum using the following assay. A monoclonal antibody raised against mammalian, recombinantly produced OP-1 using standard immunology techniques well described in the art and described generally in Example 8, was immobilized by passing the antibody over an activated agarose gel (e.g., Affi-GelTM, from Bio-Rad Laboratories, Richmond, CA, prepared following manufacturer's instructions), and used to purify OP-1 from serum. Human serum then was passed over the column and eluted with 3M K-thiocyanate. K-thiocyanate fractions then were dialyzed in 6M urea, 20mM PO₄, pH 7.0, applied to a C8 HPLC column, and eluted with a 20 minute, 25-50% acetonitrile/0.1% TFA gradient. Mature, recombinantly produced OP-1 homodimers elute between 20-22 minutes.

- 78 -

Fractions then were collected and tested for the presence of OP-1 by standard immunoblot. Fig. 7 is an immunoblot showing OP-1 in human sera under reducing and oxidized conditions. In the figure, lanes 1 and 4 are OP-1 standards, run under oxidized (lane 1) and reduced (lane 4) conditions. Lane 5 shows molecular weight markers at 17, 27 and 39 kDa. Lanes 2 and 3 are human sera OP-1, run under oxidized (lane 2) and reduced (lane 3) conditions.

10

Example 3. Morphogen-Induced CAM Expression

The morphogens described herein induce CAM expression as part of their induction of morphogenesis. CAMs are morphoregulatory molecules identified in all tissues as an essential step in tissue development. N-CAMs, which comprise at least 3 isoforms (N-CAM-180, N-CAM-140 and N-CAM-120, where "180", "140" and "120" indicate the apparent molecular weights of the isoforms as measured by polyacrylamide gel electrophoresis) are expressed at least transiently in developing tissues, and permanently in nerve tissue. Both the N-CAM-180 and N-CAM-140 isoforms are expressed in both developing and adult tissue. The N-CAM-120 isoform is found only in adult tissue. Another neural CAM is L1.

CAMs are implicated in normal tissue development; N-CAMs are implicated in appropriate neural development, including appropriate neurulation, neuronal migration, fasciculation, and synaptogenesis. Inhibition of N-CAM production, as by complexing the molecule with an N-CAM-specific antibody, inhibits retina organization, including retinal axon migration, and axon regeneration in the peripheral nervous system,

- 79 -

as well as axon synapsis with target muscle cells. CAMs also have been postulated as part of a morphoregulatory pathway whose activity is induced by a to date unidentified molecule (See, for example, 5 Edelman, G.M. (1986) Ann. Rev. Cell Biol. 2:81-116). Without being limited to any given theory, the morphogens described herein may act as the inducer of this pathway.

10 The morphogens described herein can stimulate CAM production. As described below, the morphogens stimulate L1 and N-CAM production, including all three isoforms of the N-CAM molecule, in nerve tissue.

15 In this example NG108-15 cells were cultured for 4 days in the presence of increasing concentrations of OP-1 and standard Western blots performed on whole cells extracts. The NG10875 cell line is a hybrid cell line (neuroblastoma x glioma, American Type Culture 20 Collection, Rockville, MD). N-CAM isoforms were detected with an antibody which crossreacts with all three isoforms, mAb H28.123, obtained from Sigma Chemical Co., St. Louis, the different isoforms being distinguishable by their different mobilities on an 25 electrophoresis gel. Control NG108-15 cells (untreated) express both the 140 kDa and the 180 kDa isoforms, but not the 120 kDa, as determined by western blot analyses using up to 100 μ g of protein. As shown in Fig.8, treatment of NG108-15 cells with OP-1 resulted 30 in a dose-dependent increase in the expression of the 180 kDa and 140 kDa isoforms, as well as the induction of the 120 kDa isoform. Fig. 8B is a Western blot of OP1-treated NG108-15 cell extracts, probed with mAb H28.123, showing the induction of all three isoforms.

- 80 -

Fig. 8A is a dose response curve of N-CAM-180 and N-CAM-140 induction as a function of morphogen concentration. N-CAM-120 is not shown in the graph as it could not be quantitated in control cells. However, as is clearly evident from the Western blot in Fig. 8A, N-CAM-120 is induced in response to morphogen treatment. The induction of the 120 isoform also indicates that morphogen-induced redifferentiation of transformed cells stimulates not only redifferentiation of these cells from a transformed phenotype, but also differentiation to a phenotype corresponding to a developed cell. The differential induction of N-CAM 180 and 140 isoforms seen may be because constitutive expression of the 140 isoform is close to maximum. In addition, the increase in N-CAM expression corresponded in a dose-dependent manner with the morphogen induction of multicellular aggregates.

In addition, the cell aggregation effects of OP-1 on NG108-15 cells can be inhibited with anti-N-CAM antibodies or antisense N-CAM oligonucleotides. Antisense oligonucleotides can be made synthetically on a nucleotide synthesizer, using standard means known in the art. Preferably, phosphorothioate oligonucleotides ("S-oligos") are prepared, to enhance transport of the nucleotides across cell membranes. Concentrations of both N-CAM antibodies and N-CAM antisense oligonucleotides sufficient to inhibit N-CAM induction also inhibited formation of multilayered cell aggregates. Specifically, incubation of morphogen-treated NG108-115 cells with 0.3-3 μ M N-CAM antisense S-oligos, 5-500 μ M unmodified N-CAM antisense oligos,

- 81 -

or 10 μ g/ml mAb H28.123 significantly inhibits cell aggregation. It is likely that morphogen treatment also stimulates other CAMs, as inhibition is not complete.

5

The experiments also have been performed with soluble morphogen (e.g., mature OP-1 associated with its pro domain) which also specifically induced CAM expression.

10

Example 4. Effect of Morphogen Neutralization on Embryogenesis

As described in Example 7, below, at least one
15 morphogen, OP2, is found principally in early developing embryos (8-day embryos). As described below, morphogen neutralization with morphogen-specific antibodies inhibits embryogenesis.

20 Morphogen inhibition in developing embryos inhibits tissue and organ development. Specifically, 9-day mouse embryo cells, cultured in vitro under standard culturing conditions, were incubated in the presence and absence of an OP-1-specific monoclonal antibody
25 prepared using recombinantly produced, purified mature OP-1 as the immunogen. The antibody was prepared using standard antibody production means well known in the art and essentially as described for Example 9, below. After two days, the effect of the antibody on the
30 developing embryo was evaluated by histology using standard histology procedures well known in the art. As determined by histological examination, the OP-1-specific antibody specifically inhibits eye lobe formation in the developing embryo. In particular, the

- 82 -

diencephalon outgrowth does not develop. In addition, the heart is malformed and enlarged. Moreover, in separate immunolocalization studies on embryo sections with labelled OP-1 specific antibody, the OP-1-specific antibody localizes to neural epithelia.

Similarly, morphogen activity may be demonstrated in fetal development in the mouse model using the following assay. Single lip injections comprising 10-100 μ g/injection of morphogen-specific antibody are administered to pregnant female mice during each day of the gestation period and tissue development (e.g., bone development) in treated and control new mice evaluated by standard histomorphometric analysis at birth.

Finally, stimulation of endogenous morphogen antibody production in egg-laying hens interferes with shell formation in the developing eggs.

All of these data demonstrate that inhibition of morphogen activity significantly interferes with tissue development during embryogenesis.

Example 5. Effect of Morphogen Neutralization on Juvenile Tissue Development

The effect of the morphogens described herein on tissue development in developing mammals also may be demonstrated using neutralizing antibodies specific for particular morphogens and assessing the effect of these antibodies on tissue development as described below. Specifically, anti-morphogen monoclonal and/or polyclonal antibodies may be prepared using standard methodologies including, for example, the protocol

- 83 -

provided in Example 8, below, and provided to juveniles to inhibit the activity of endogenous morphogens.

Generally, purified antibodies are provided
5 regularly to new born mice, e.g., 10-
100 μ g/injection/day for 10-15 days. At 10 or 21 days,
the mice are sacrificed and the effect of morphogen on
bone development assessed by body weight, gross visual
examination and histology. In this example, anti-OP-1
10 antibodies were used in 10 μ g injections/day for 14
days, and the mice were sacrificed at 21 days. As is
dramatically demonstrated in Fig. 9, mice treated with
OP-1 specific antibody show consistent and significant
stunted growth, including reduced body length and body
15 weight, (9B) as compared with untreated mice (9A).
Histological examination showed reduced bone growth as
evidenced by reduced bone size in the treated mice.

In a variation on this protocol, single lip
20 injections also may be provided to older juveniles and
adult mice (e.g., 10-100 μ g) over a prolonged time
(e.g., 10-15 days) to evaluate the effect of morphogen
neutralization on bone growth and bone integrity and to
evaluate the onset of osteoporosis.

25

Example 6. Morphogen Treatment of Osteoporosis

6.1 Effect of Morphogen on Trabecular Bone in Ovariectomized (OVX) Rats

30

Aged individuals, and particularly postmenopausal
women are particularly at risk for osteoporosis.
Provided below is an animal osteoporosis model
demonstrating the ability of morphogens to substantially

- 84 -

inhibit and/or reduce the tissue damage effects associated with osteoporosis, wherein osteoporosis is induced by ovary removal in rats. Bone growth is evaluated in these animals by measuring serum alkaline phosphatase and osteocalcin levels in treated and untreated rats.

Forty Long-Evans rats (Charles River Laboratories, Wilmington) weighing about 200g each are ovariectomized (OVX) using standard surgical procedures, and ten rats are sham-operated. The ovariectomization of the rats produces an osteoporotic condition within the rats as a result of decreased estrogen production. Food and water are provided ad libitum. Eight days after ovariectomy, the rats, prepared as described above, were divided into five groups: (A), 10 sham-operated rats; (B), 10 ovariectomized rats receiving 1 ml of phosphate-buffered saline (PBS) i.v. in the tail vein; (C) 10 ovariectomized rats receiving about 1 mg of $17\beta E_2$ (17β -estradiol E_2) by intravenous injection through the tail vein; (D) 9 ovariectomized rats receiving daily injections of approximately $2\mu g$ of morphogen by tail vein for 22 days; and (E) 9 ovariectomized rats receiving daily injections of approximately $20\mu g$ of morphogen by tail vein for 22 days. In this example, OP-1 was the morphogen tested.

On the 15th and 21st day of the study, each rat was injected with 5 mg of tetracycline, and on day 22, the rats were sacrificed. The body weights, uterine

- 85 -

weights, serum alkaline phosphate levels, serum calcium levels and serum osteocalcin levels then were determined for each rat. The results are shown in Tables III and IV.

5

Table IIIBody Weights, Uterine Weights and Alkaline Phosphatase

Group	<u>Body Weights</u>	<u>Uterine Weights</u>	<u>Alk. Phosphatase</u>
	(g)	(g)	(U/L)
10 A-SHAM	250.90 \pm 17.04	0.4192 \pm 0.10	43.25 \pm 6.11
B-OVX+PBS	273.40 \pm 16.81	0.1650 \pm 0.04	56.22 \pm 6.21
C-OVX+E2	241.66 \pm 21.54	0.3081 \pm 0.03	62.66 \pm 4.11
D-OVX+OP-1	266.67 \pm 10.43	0.1416 \pm 0.03	58.09 \pm 12.97
15 (2 μ g)			
E-OVX+OP-1	272.40 \pm 20.48	0.1481 \pm 0.05	66.24 \pm 15.74
(20 μ g)			

TABLE IV

20

Serum Calcium and Serum Osteocalcin Levels

Group	<u>Serum Calcium</u>	<u>Serum Osteocalcin</u>
	(ng/dl)	(ng/ml)
25 A-SHAM	8.82 \pm 1.65	64.66 \pm 14.77
B-OVX+PBS	8.95 \pm 1.25	69.01 \pm 10.20
C-OVX+E2	9.20 \pm 1.39	67.13 \pm 17.33
D-OVX+OP-1	8.77 \pm 0.95	148.50 \pm 84.11
30 (2 μ g)		
E-OVX+OP-1	8.67 \pm 1.94	182.42 \pm 52.11
(20 μ g)		

- 86 -

The results presented in Table III and IV show that intravenous injection of morphogen into ovariectomized rats produces a significant increase in serum alkaline phosphatase and serum osteocalcin levels and demonstrates that systemic administration of the morphogen stimulates bone formation in osteoporotic bone.

6.2 Histomorphometric Analysis of Morphogen on the
Tibia Diaphysis in Ovariectomized(OVX) Rats

Fifteen female Long-Evans rats weighing about 160 g were ovariectomized (OVX) to produce an osteoporotic condition and five rats were sham operated (Charles River Laboratories, Wilmington, MA.) as described for Example 8. Food and water were provided ad libitum. Twenty-two days after ovariectomy, the rats were divided into four groups: (A) sham-operated (1 ml of PBS by intravenous injection through tail vein (5 rats); (B) OVX, into which nothing was injected (5 rats); (C) OVX, receiving about 1 mg of $17\beta E_2$ by intravenous injection through the tail vein (5 rats), and (D) OVX, receiving about 1 μg of morphogen by intravenous injection through the tail vein (5 rats). In this example, OP-1 was morphogen tested.

The rats were injected daily as described for seven days, except no injections were given on the thirteenth day. The rats then were sacrificed on the nineteenth day. The tibial diaphyseal long bones then were removed and fixed in ethanol and histomorphometric analysis was carried out using standard procedures well known in the art. The results are shown in Table V.

- 87 -

Table V

MEASUREMENT	(A) CONTROL	(B) OVX	(C) OVX + E ₂	(D) OVX + OP-1
5 Longitudinal Growth Rate ($\mu\text{m}/\text{day}$)	20.2 \pm 0.3	19.4 \pm 0.2	4.9 \pm 0.5	17.9 \pm 0.9
Cancellous Bone Volume (BV/TV, 10 bone vol/total vol)	20.2 \pm 1.5	13.0 \pm 1.6	13.7 \pm 2.1	16.6 \pm 1.8
Cancellous Bone Perimeter (mm)	16.2 \pm 1.8	9.6 \pm 0.9	11.5 \pm 1.1	12.2 \pm 0.7
15 Labeled Cancellous Perimeter (%)	35.5 \pm 1.5	51.9 \pm 5.6	58.0 \pm 4.2	39.2 \pm 1.9
Mineral Apposition Rate ($\mu\text{m}/\text{day}$)	1.76 \pm 0.14	2.25 \pm 0.16	1.87 \pm 0.08	1.86 \pm 0.20

20

The results presented in Table V confirm the results of Example 6.1, namely that intravenous injection of OP-1 into ovariectomized rats stimulates bone growth for bone which had been lost due to the drop in estrogen within the individual rat. Specifically, the inhibition of cancellous bone volume in OVX rats is repaired by the systemically provided morphogen. In addition, in morphogen-treated rats the labelled cancellous perimeter and mineral apposition rate now return to levels measured in the control, sham-operated rats. Moreover, morphogen treatment does not inhibit longitudinal bone growth, unlike estrogen

30

- 88 -

treatment, which appears to inhibit bone growth significantly. Accordingly, systemic administration of a morphogen in therapeutically effective concentrations effectively inhibits loss of bone mass in a mammal
5 without inhibiting natural bone formation.

Example 7. Identification of Morphogen-Expressing
Tissue

10 Determining the tissue distribution of morphogens may be used to identify different morphogens expressed in a given tissue, as well as to identify new, related morphogens. Tissue distribution also may be used to
15 identify useful morphogen-producing tissue for use in screening and identifying candidate morphogen-stimulating agents. The morphogens (or their mRNA transcripts) readily are identified in different tissues using standard methodologies and minor
20 modifications thereof in tissues where expression may be low. For example, protein distribution may be determined using standard Western blot analysis or immunofluorescent techniques, and antibodies specific to the morphogen or morphogens of interest. Similarly,
25 the distribution of morphogen transcripts may be determined using standard Northern hybridization protocols and transcript-specific probes.

Any probe capable of hybridizing specifically to a transcript, and distinguishing the transcript of
30 interest from other, related transcripts may be used. Because the morphogens described herein share such high sequence homology in their active, C-terminal domains, the tissue distribution of a specific morphogen transcript may best be determined using a probe

- 89 -

specific for the pro region of the immature protein and/or the N-terminal region of the mature protein. Another useful sequence is the 3' non-coding region flanking and immediately following the stop codon.

- 5 These portions of the sequence vary substantially among the morphogens of this invention, and accordingly, are specific for each protein. For example, a particularly useful Vgr-1-specific probe sequence is the PvuII-SacI fragment, a 265 bp fragment encoding both a portion of
- 10 the untranslated pro region and the N-terminus of the mature sequence (see Lyons et al. (1989) PNAS 86:4554-4558 for a description of the cDNA sequence). Similarly, particularly useful mOP-1-specific probe sequences are the BstXI-BglI fragment, a 0.68 Kb
- 15 sequence that covers approximately two-thirds of the mOP-1 pro region; a StuI-StuI fragment, a 0.2 Kb sequence immediately upstream of the 7-cysteine domain; and the Earl-PstI fragment, an 0.3 Kb fragment containing a portion of the 3'untranslated sequence
- 20 (See Seq. ID No. 18, where the pro region is defined essentially by residues 30-291.) Similar approaches may be used, for example, with hOP-1 (Seq. ID No. 16) or human or mouse OP-2 (Seq. ID Nos. 20 and 22.)

- 25 Using these morphogen-specific probes, which may be synthetically engineered or obtained from cloned sequences, morphogen transcripts can be identified in mammalian tissue, using standard methodologies well known to those having ordinary skill in the art.
- 30 Briefly, total RNA is prepared from various adult murine tissues (e.g., liver, kidney, testis, heart, brain, thymus and stomach) by a standard methodology such as by the method of Chomczyaski et al. ((1987) Anal. Biochem 162:156-159) and described below. Poly

- 90 -

- (A)+ RNA is prepared by using oligo (dT)-cellulose chromatography (e.g., Type 7, from Pharmacia LKB Biotechnology, Inc.). Poly (A)+ RNA (generally 15 μ g) from each tissue is fractionated on a 1% agarose/formaldehyde gel and transferred onto a Nytran membrane (Schleicher & Schuell). Following the transfer, the membrane is baked at 80°C and the RNA is cross-linked under UV light (generally 30 seconds at 1 mW/cm²). Prior to hybridization, the appropriate probe is denatured by heating. The hybridization is carried out in a lucite cylinder rotating in a roller bottle apparatus at approximately 1 rev/min for approximately 15 hours at 37°C using a hybridization mix of 40% formamide, 5 x Denhardts, 5 x SSPE, and 0.1% SDS. Following hybridization, the non-specific counts are washed off the filters in 0.1 x SSPE, 0.1% SDS at 50°C.

Examples demonstrating the tissue distribution of various morphogens, including Vgr-1, OP-1, BMP2, BMP3, BMP4, BMP5, GDF-1, and OP-2 in developing and adult tissue are disclosed in international application US92/01968 (WO92/15323), and in Ozkaynak, et al., (1991) Biochem. Biophys. Res. Commun. 179:116-123, and Ozkaynak, et al. (1992) (J. Biol. Chem. 267: 25220-25227), the disclosures of which are incorporated herein by reference. Using the general probing methodology described herein, northern blot hybridizations using probes specific for these morphogens to probe brain, spleen, lung, heart, liver and kidney tissue indicate that kidney-related tissue appears to be the primary expression source for OP-1, with brain, heart and lung tissues being secondary sources. Lung tissue appears to be the primary tissue expression source for Vgr-1, BMP5, BMP4 and BMP3. Lower

- 91 -

levels of Vgr-1 also are seen in kidney and heart tissue, while the liver appears to be a secondary expression source for BMP5, and the spleen appears to be a secondary expression source for BMP4. GDF-1
5 appears to be expressed primarily in brain tissue.

Of particular relevance to the present application, OP-1 also is detected in adult rat stomach and gut tissue. Moreover, OP-2 appears to be expressed
10 primarily in early embryonic tissue. Specifically, northern blots of murine embryos and 6-day post-natal animals shows abundant OP2 expression in 8-day embryos. Expression is reduced significantly in 17-day embryos and is not detected in post-natal animals.

15

In addition, labelled soluble OP-1 (iodinated with ^{125}I , using standard labelling procedures well known in the art) and injected into the rat tail vein also is localized to the stomach tissue within 30 minutes of
20 injection.

Example 8. Detecting Morphogenic Protein in Solution
by Immunoassay

25 Morphogens are readily detected in solution with a standard immunoassay, using a polyclonal or monoclonal antibody specific for that protein and standard Western blot, ELISA (enzyme-linked immunoabsorbant assay) or other immunoassay technique well known in the art. A
30 currently preferred, exemplary protocol for an ELISA assay, as well as means for generating morphogen-specific antibody are presented below. Standard protocols for antibody production, Western blot and other immunoassays also are described, for example, in

- 92 -

Molecular Cloning A Laboratory Manual, Sambrook et al., eds. 1989, Cold Spring Harbor Press, Cold Spring Harbor, NY. Standard ELISA technique is described, for example, by Engvall (1980) Methods Enzymol. 70:419-439.

5

8.1 Morphogen-Specific Antiserum

Polyclonal antibody was prepared as follows. Each rabbit was given a primary immunization of 100 ug/500
10 μ l E. coli-produced OP-1 monomer (amino acids 328-431 in SEQ ID NO:5) in 0.1% SDS mixed with 500 μ l

Complete Freund's Adjuvant. The antigen was injected subcutaneously at multiple sites on the back and flanks of the animal. The rabbit was boosted after a month in
15 the same manner using incomplete Freund's Adjuvant. Test bleeds are taken from the ear vein seven days later. Two additional boosts and test bleeds were performed at monthly intervals until antibody against OP-1 was detected in the serum using an ELISA assay.
20 Then, the rabbit was boosted monthly with 100 μ g of antigen and bled (15 ml per bleed) at days seven and ten after boosting.

8.2 Morphogen-Specific Antibody

25

Monoclonal antibody specific for a given morphogen was prepared as follows. A mouse was given two injections of E. coli produced OP-1 monomer. The first injection contains 100 μ g of OP-1 in complete Freund's
30 adjuvant and was given subcutaneously. The second injection contained 50 μ g of OP-1 in incomplete adjuvant and was given intraperitoneally. The mouse then received a total of 230 μ g of OP-1 (amino acids 307-431 in SEQ ID NO:5) in four intraperitoneal

- 93 -

injections at various times over an eight month period. One week prior to fusion, both mice were boosted intraperitoneally with 100 μ g of OP-1 (307-431) and 30 μ g of the N-terminal peptide (Ser293-Asn309-Cys) conjugated through the added cysteine to bovine serum albumin with SMCC crosslinking agent. This boost was repeated five days (IP), four days (IP), three days (IP) and one day (IV) prior to fusion. The mouse spleen cells then were fused to commercially available myeloma cells at a ratio of 1:1 using PEG 1500 (Boeringer Mannheim, Germany), and the cell fusion plated and screened for OP-1-specific antibodies using OP-1 (307-431) as antigen. The cell fusion and monoclonal screening then were according to standard procedures well described in standard texts widely available in the art e.g., Maniatis et al. Molecular Cloning A Laboratory Manual, Cold Spring Harbor Press.

8.3 Morphogen ELISA

1 μ g/100 μ l of affinity-purified polyclonal rabbit IgG specific for OP-1 was added to each well of a 96-well plate and incubated at 37°C for an hour. The wells were washed four times with 0.167M sodium borate buffer with 0.15 M NaCl (BSB), pH 8.2, containing 0.1% Tween 20. To minimize non-specific binding, the wells are blocked by filling completely with 1% bovine serum albumin (BSA) in BSB and incubating for 1 hour at 37°C. The wells are then washed four times with BSB containing 0.1% Tween 20. A 100 μ l aliquot of an appropriate dilution of each of the test samples of cell culture supernatant was added to each well in triplicate and incubated at 37°C for 30 min. After incubation, 100 μ l biotinylated rabbit anti-OP-1 serum

- 94 -

(stock solution is about 1 mg/ml and diluted 1:400 in BSB containing 1% BSA before use) are added to each well and incubated at 37°C for 30 min. The wells were then washed four times with BSB containing 0.1% Tween

5 20. 100 µl strepavidin-alkaline (Southern Biotechnology Associates, Inc. Birmingham, Alabama, diluted 1:2000 in BSB containing 0.1% Tween 20 before use) was added to each well and incubated at 37°C for 30 min. The plates were washed four times with 0.5M

10 Tris buffered Saline (TBS), pH 7.2. 50µl substrate (ELISA Amplification System Kit, Life Technologies, Inc., Bethesda, MD) was added to each well incubated at room temperature for 15 min. Then, 50 µl amplifier (from the same amplification system kit) is added and

15 incubated for another 15 min at room temperature. The reaction was stopped by the addition of 50 µl 0.3 M sulphuric acid. The OD at 490 nm of the solution in each well was recorded. To quantitate OP-1 in culture media, an OP-1 standard curve was performed in parallel

20 with the test samples.

The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The present embodiments are

25 therefore to be considered in all respects as illustrative and not restrictive, the scope of the invention being indicated by the appended claims rather than by the foregoing description, and all changes which come within the meaning and range of equivalency

30 of the claims are therefore intended to be embraced therein.

- 95 -

SEQUENCE LISTING

5 (1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: CREATIVE BIOMOLECULES, INC.
- (B) STREET: 35 SOUTH STREET
- 10 (C) CITY: HOPKINTON
- (D) STATE: MA
- (E) COUNTRY: USA
- (F) POSTAL CODE (ZIP): 01748
- (G) TELEPHONE: 1-508-435-9001
- 15 (H) TELEFAX: 1-508-435-0454
- (I) TELEX:

(ii) TITLE OF INVENTION: MORPHOGEN-ENRICHED DIETARY COMPOSITION

20 (iii) NUMBER OF SEQUENCES: 33

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: CREATIVE BIOMOLECULES, INC.
- (B) STREET: 35 SOUTH STREET
- 25 (C) CITY: HOPKINTON
- (D) STATE: MA
- (E) COUNTRY: USA
- (F) ZIP: 01748

30 (v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- 35 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: KELLEY, ROBIN D.
- (B) REGISTRATION NUMBER: 34,637
- 40 (C) REFERENCE/DOCKET NUMBER: CRP-071

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 617/248-7000
- (B) TELEFAX: 617/248-7100

45 (2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids
- (B) TYPE: amino acid
- 50 (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 96 -

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

5

(A) NAME/KEY: Protein

(B) LOCATION: 1..97

(D) OTHER INFORMATION: /label= GENERIC-SEQ1

10

/note= "WHEREIN EACH XAA INDEPENDENTLY INDICATES
ONE OF THE 20 NATURALLY-OCCURRING L-ISOMER, A-AMINO
ACIDS, OR A DERIVATIVE THEREOF."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10 15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Cys Xaa Xaa Xaa
20 25 30

20

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
35 40 45

25

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Cys Xaa Xaa
50 55 60

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
65 70 75 80

30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Cys
85 90 95

Xaa

35

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

40

(A) LENGTH: 97 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

45

- 97 -

(ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..97

(D) OTHER INFORMATION: /label= GENERIC-SEQ2

5 /note= "WHEREIN EACH XAA INDEPENDENTLY INDICATES
ONE OF THE 20 NATURALLY OCCURRING L-ISOMER A-AMINO
ACIDS, OR A DERIVATIVE THEREOF."

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10 15
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Cys Xaa Xaa Xaa
20 25 30
Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
35 40 45
20 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Cys Xaa Xaa
50 55 60
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
25 65 70 75 80
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Cys
85 90 95
30 Xaa

(2) INFORMATION FOR SEQ ID NO:3:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 97 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: protein

- 98 -

(ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..97

(D) OTHER INFORMATION: /label= GENERIC-SEQ3

5 /note= "WHEREIN EACH XAA IS INDEPENDENTLY SELECTED
FROM A GROUP OF ONE OR MORE SPECIFIED AMINO ACIDS
AS DEFINED IN THE SPECIFICATION."

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Leu Tyr Val Xaa Phe Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa Xaa Ala
1 5 10 15
Pro Xaa Gly Xaa Xaa Ala Xaa Tyr Cys Xaa Gly Xaa Cys Xaa Xaa Pro
20 25 30
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala Xaa Xaa Xaa Xaa Leu
35 40 45
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Cys Xaa Pro
50 55 60
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
25 65 70 75 80
Val Xaa Leu Xaa Xaa Xaa Xaa Xaa Met Xaa Val Xaa Xaa Cys Gly Cys
85 90 95
30 Xaa

(2) INFORMATION FOR SEQ ID NO:4:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 102 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: protein

- 99 -

(ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..102

(D) OTHER INFORMATION: /label= GENERIC-SEQ4

5 /note= "WHEREIN EACH XAA IS INDEPENDENTLY SELECTED
FROM A GROUP OF ONE OR MORE SPECIFIED AMINO ACIDS
AS DEFINED IN THE SPECIFICATION."

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Cys Xaa Xaa Xaa Xaa Leu Tyr Val Xaa Phe Xaa Xaa Xaa Gly Trp Xaa
1 5 10 15

15 Xaa Trp Xaa Xaa Ala Pro Xaa Gly Xaa Xaa Ala Xaa Tyr Cys Xaa Gly
 20 25 30

Xaa Cys Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala
 35 40 45

20 Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 50 55 60

25 Xaa Cys Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa
65 70 75 80

Xaa Xaa Xaa Xaa Xaa Val Xaa Leu Xaa Xaa Xaa Xaa Xaa Met Xaa Val
 85 90 95

30 Xaa Xaa Cys Gly Cys Xaa
 100

(2) INFORMATION FOR SEQ ID NO:5:

35 (1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 139 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

45 (F) TISSUE TYPE: HIPPOCAMPUS

- 100 -

(ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..139

(D) OTHER INFORMATION: /label= hOP1-MATURE

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

10 Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser Lys Thr Pro Lys
 1 5 10 15
 Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu Asn Ser Ser Ser
 20 25 30
 15 Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg
 35 40 45
 Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala
 50 55 60
 20 Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met Asn
 65 70 75 80
 Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn Pro
 25 85 90 95
 Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala Ile
 100 105 110
 30 Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr
 115 120 125
 Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 130 135

35

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 139 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: protein

45

(vi) ORIGINAL SOURCE:

(A) ORGANISM: MURIDAE

(F) TISSUE TYPE: EMBRYO

- 101 -

(ix) FEATURE:

- (A) NAME/KEY: Protein
 (B) LOCATION: 1..139
 (D) OTHER INFORMATION: /label= NOP1-MATURE

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

10 Ser Thr Gly Gly Lys Gln Arg Ser Gln Asn Arg Ser Lys Thr Pro Lys
 1 5 10 15
 Asn Gln Glu Ala Leu Arg Met Ala Ser Val Ala Glu Asn Ser Ser Ser
 20 25 30
 15 Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg
 35 40 45
 Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala
 50 55 60
 20 Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met Asn
 65 70 75 80
 Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn Pro
 85 90 95
 25 Asp Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala Ile
 100 105 110
 30 Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr
 115 120 125
 Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 130 135

35

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 139 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

45

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: HOMO SAPIENS
 (F) TISSUE TYPE: HIPPOCAMPUS

- 102 -

(ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..139

(D) OTHER INFORMATION: /label= HOP2-MATURE

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

10 Ala Val Arg Pro Leu Arg Arg Arg Gln Pro Lys Lys Ser Asn Glu Leu
 1 5 10 15
 Pro Gln Ala Asn Arg Leu Pro Gly Ile Phe Asp Asp Val His Gly Ser
 20 25 30
 15 His Gly Arg Gln Val Cys Arg Arg His Glu Leu Tyr Val Ser Phe Gln
 35 40 45
 Asp Leu Gly Trp Leu Asp Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala
 50 55 60
 20 Tyr Tyr Cys Glu Gly Glu Cys Ser Phe Pro Leu Asp Ser Cys Met Asn
 65 70 75 80
 Ala Thr Asn His Ala Ile Leu Gln Ser Leu Val His Leu Met Lys Pro
 85 90 95
 25 Asn Ala Val Pro Lys Ala Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr
 100 105 110
 30 Ser Val Leu Tyr Tyr Asp Ser Ser Asn Asn Val Ile Leu Arg Lys His
 115 120 125
 Arg Asn Met Val Val Lys Ala Cys Gly Cys His
 130 135

35

-(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 139 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: protein

45

(vi) ORIGINAL SOURCE:

(A) ORGANISM: MURIDAE

(F) TISSUE TYPE: EMBRYO

- 103 -

(ix) FEATURE:

- (A) NAME/KEY: Protein
(B) LOCATION: 1..139
(D) OTHER INFORMATION: /label= MOP2-MATURE

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

10 Ala Ala Arg Pro Leu Lys Arg Arg Gln Pro Lys Lys Thr Asn Glu Leu
1 5 10 15
Pro His Pro Asn Lys Leu Pro Gly Ile Phe Asp Asp Gly His Gly Ser
20 25 30
15 Arg Gly Arg Glu Val Cys Arg Arg His Glu Leu Tyr Val Ser Phe Arg
35 40 45
20 Asp Leu Gly Trp Leu Asp Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala
50 55 60
Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asp Ser Cys Met Asn
65 70 75 80
25 Ala Thr Asn His Ala Ile Leu Gln Ser Leu Val His Leu Met Lys Pro
85 90 95
Asp Val Val Pro Lys Ala Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr
100 105 110
30 Ser Val Leu Tyr Tyr Asp Ser Ser Asn Asn Val Ile Leu Arg Lys His
115 120 125
Arg Asn Met Val Val Lys Ala Cys Gly Cys His
130 135

35

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 101 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

45

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: bovinæ

- 104 -

(ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..101

(D) OTHER INFORMATION: /label= CBMP-2A-FX

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

10 Cys Lys Arg His Pro Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asn
 1 5 10 15
 Asp Trp Ile Val Ala Pro Pro Gly Tyr His Ala Phe Tyr Cys His Gly
 20 25 30
 15 Glu Cys Pro Phe Pro Leu Ala Asp His Leu Asn Ser Thr Asn His Ala
 35 40 45
 Ile Val Gln Thr Leu Val Asn Ser Val Asn Ser Lys Ile Pro Lys Ala
 50 55 60
 20 Cys Cys Val Pro Thr Glu Leu Ser Ala Ile Ser Met Leu Tyr Leu Asp
 65 70 75 80
 25 Glu Asn Glu Lys Val Val Leu Lys Asn Tyr Gln Asp Met Val Val Glu
 85 90 95
 Gly Cys Gly Cys Arg
 100

30 (2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 101 amino acids

(B) TYPE: amino acid

35 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

40 (vi) ORIGINAL SOURCE:

(A) ORGANISM: HOMO SAPIENS

(F) TISSUE TYPE: hippocampus

45 (ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..101

(D) OTHER INFORMATION: /label= CBMP-2B-FX

- 105 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

5 Cys Arg Arg His Ser Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asn
 1 5 10 15
 Asp Trp Ile Val Ala Pro Pro Gly Tyr Gln Ala Phe Tyr Cys His Gly
 20 25 30
 10 Asp Cys Pro Phe Pro Leu Ala Asp His Leu Asn Ser Thr Asn His Ala
 35 40 45
 Ile Val Gln Thr Leu Val Asn Ser Val Asn Ser Ser Ile Pro Lys Ala
 50 55 60
 15 Cys Cys Val Pro Thr Glu Leu Ser Ala Ile Ser Met Leu Tyr Leu Asp
 65 70 75 80
 Glu Tyr Asp Lys Val Val Leu Lys Asn Tyr Gln Glu Met Val Val Glu
 85 90 95
 20 Gly Cys Gly Cys Arg
 100

(2) INFORMATION FOR SEQ ID NO:11:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 102 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 30 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein
 (vi) ORIGINAL SOURCE:
 35 (A) ORGANISM: DROSOPHILA MELANOGASTER
 (ix) FEATURE:
 (A) NAME/KEY: Protein
 (B) LOCATION: 1..101
 40 (D) OTHER INFORMATION: /label= DPP-FX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

45 Cys Arg Arg His Ser Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asp
 1 5 10 15
 Asp Trp Ile Val Ala Pro Leu Gly Tyr Asp Ala Tyr Tyr Cys His Gly
 20 25 30
 50

- 106 -

Lys Cys Pro Phe Pro Leu Ala Asp His Phe Asn Ser Thr Asn His Ala
 35 40 45
 Val Val Gln Thr Leu Val Asn Asn Asn Asn Pro Gly Lys Val Pro Lys
 5 50 55 60
 Ala Cys Cys Val Pro Thr Gln Leu Asp Ser Val Ala Met Leu Tyr Leu
 65 70 75 80
 10 Asn Asp Gln Ser Thr Val Val Leu Lys Asn Tyr Gln Glu Met Thr Val
 85 90 95
 Val Gly Cys Gly Cys Arg
 100

15

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 102 amino acids
 20 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

 (ii) MOLECULE TYPE: protein
 25
 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: XENOPUS

 (ix) FEATURE:
 30 (A) NAME/KEY: Protein
 (B) LOCATION: 1..102
 (D) OTHER INFORMATION: /label= VGL-FX

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Cys Lys Lys Arg His Leu Tyr Val Glu Phe Lys Asp Val Gly Trp Gln
 1 5 10 15
 40 Asn Trp Val Ile Ala Pro Gln Gly Tyr Met Ala Asn Tyr Cys Tyr Gly
 20 25 30
 Glu Cys Pro Tyr Pro Leu Thr Glu Ile Leu Asn Gly Ser Asn His Ala
 35 40 45
 45 Ile Leu Gln Thr Leu Val His Ser Ile Glu Pro Glu Asp Ile Pro Leu
 50 55 60
 50 Pro Cys Cys Val Pro Thr Lys Met Ser Pro Ile Ser Met Leu Phe Tyr
 65 70 75 80

- 107 -

Asp Asn Asn Asp Asn Val Val Leu Arg His Tyr Glu Asn Met Ala Val
 85 90 95

5 Asp Glu Cys Gly Cys Arg
 100

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 102 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

(A) ORGANISM: MURIDAE

20 (ix) FEATURE:

- (A) NAME/KEY: Protein
 (B) LOCATION: 1..102
 (D) OTHER INFORMATION: /label= VGR-1-FX

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Cys Lys Lys His Glu Leu Tyr Val Ser Phe Gln Asp Val Gly Trp Gln
 1 5 10 15

30 Asp Trp Ile Ile Ala Pro Lys Gly Tyr Ala Ala Asn Tyr Cys Asp Gly
 20 25 30

Glu Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His Ala
 35 35 40 45

Ile Val Gln Thr Leu Val His Val Met Asn Pro Glu Tyr Val Pro Lys
 50 55 60

40 Pro Cys Cys Ala Pro Thr Lys Val Asn Ala Ile Ser Val Leu Tyr Phe
 65 70 75 80

Asp Asp Asn Ser Asn Val Ile Leu Lys Lys Tyr Arg Asn Met Val Val
 85 90 95

45 Arg Ala Cys Gly Cys His
 100

- 108 -

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 106 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
 (F) TISSUE TYPE: brain

(ix) FEATURE:
 (A) NAME/KEY: Protein
 (B) LOCATION: 1..106
 (D) OTHER INFORMATION: /note= "GDF-1 (fx)"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Cys	Arg	Ala	Arg	Arg	Leu	Tyr	Val	Ser	Phe	Arg	Glu	Val	Gly	Trp	His
1				5					10					15	
Arg	Trp	Val	Ile	Ala	Pro	Arg	Gly	Phe	Leu	Ala	Asn	Tyr	Cys	Gln	Gly
		20					25						30		
Gln	Cys	Ala	Leu	Pro	Val	Ala	Leu	Ser	Gly	Ser	Gly	Gly	Pro	Pro	Ala
		35					40					45			
Leu	Asn	His	Ala	Val	Leu	Arg	Ala	Leu	Met	His	Ala	Ala	Ala	Pro	Gly
	50					55				60					
Ala	Ala	Asp	Leu	Pro	Cys	Cys	Val	Pro	Ala	Arg	Leu	Ser	Pro	Ile	Ser
65					70				75					80	
Val	Leu	Phe	Phe	Asp	Asn	Ser	Asp	Asn	Val	Val	Leu	Arg	Gln	Tyr	Glu
			85					90					95		
Asp	Met	Val	Val	Asp	Glu	Cys	Gly	Cys	Arg						
			100					105							

- 109 -

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

15 Cys Xaa Xaa Xaa Xaa
1 5

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 1822 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

30

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: HOMO SAPIENS
(F) TISSUE TYPE: HIPPOCAMPUS

35

(ix) FEATURE:

- 40 (A) NAME/KEY: CDS
(B) LOCATION: 49..1341
(C) IDENTIFICATION METHOD: experimental
(D) OTHER INFORMATION: /function= "OSTEOGENIC PROTEIN"
/product= "OP1"
/evidence= EXPERIMENTAL
/standard_name= "OP1"

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

- 110 -

	GGTGCGGGCC CGGAGCCCGG AGCCCGGGTA GCGCGTAGAG CCGGCGCG ATG CAC GTG	57
	Met His Val	
	1	
5	CGC TCA CTG CGA GCT GCG GCG CCG CAC AGC TTC GTG GCG CTC TGG GCA	105
	Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala Leu Trp Ala	
	5 10 15	
10	CCC CTG TTC CTG CTG CGC TCC GCC CTG GCC GAC TTC AGC CTG GAC AAC	153
	Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser Leu Asp Asn	
	20 25 30 35	
15	GAG GTG CAC TCG AGC TTC ATC CAC CGG CGC CTC CGC AGC CAG GAG CGG	201
	Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser Gln Glu Arg	
	40 45 50	
20	CGG GAG ATG CAG CGC GAG ATC CTC TCC ATT TTG GGC TTG CCC CAC CGC	249
	Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu Pro His Arg	
	55 60 65	
20	CCG CGC CCG CAC CTC CAG GGC AAG CAC AAC TCG GCA CCC ATG TTC ATG	297
	Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro Met Phe Met	
	70 75 80	
25	CTG GAC CTG TAC AAC GCC ATG GCG GTG GAG GAG GGC GGC GGG CCC GGC	345
	Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly Gly Pro Gly	
	85 90 95	
30	GGC CAG GGC TTC TCC TAC CCC TAC AAG GCC GTC TTC AGT ACC CAG GGC	393
	Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser Thr Gln Gly	
	100 105 110 115	
35	CCC CCT CTG GCC AGC CTG CAA GAT AGC CAT TTC CTC ACC GAC GCC GAC	441
	Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr Asp Ala Asp	
	120 125 130	
40	ATG GTC ATG AGC TTC GTC AAC CTC GTG GAA CAT GAC AAG GAA TTC TTC	489
	Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys Glu Phe Phe	
	135 140 145	
40	CAC CCA CGC TAC CAC CAT CGA GAG TTC CGG TTT GAT CTT TCC AAG ATC	537
	His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu Ser Lys Ile	
	150 155 160	
45	CCA GAA GGG GAA GCT GTC ACG GCA GCC GAA TTC CGG ATC TAC AAG GAC	585
	Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile Tyr Lys Asp	
	165 170 175	

- 111 -

	TAC	ATC	CGG	GAA	CGC	TTC	GAC	AAT	GAG	ACG	TTC	CGG	ATC	AGC	GTT	TAT	633
	Tyr	Ile	Arg	Glu	Arg	Phe	Asp	Asn	Glu	Thr	Phe	Arg	Ile	Ser	Val	Tyr	
	180					185					190					195	
5	CAG	GTG	CTC	CAG	GAG	CAC	TTG	GGC	AGG	GAA	TCG	GAT	CTC	TTC	CTG	CTC	681
	Gln	Val	Leu	Gln	Glu	His	Leu	Gly	Arg	Glu	Ser	Asp	Leu	Phe	Leu	Leu	
					200					205					210		
10	GAC	AGC	CGT	ACC	CTC	TGG	GCC	TCG	GAG	GAG	GGC	TGG	CTG	GTG	TTT	GAC	729
	Asp	Ser	Arg	Thr	Leu	Trp	Ala	Ser	Glu	Glu	Gly	Trp	Leu	Val	Phe	Asp	
				215					220					225			
15	ATC	ACA	GCC	ACC	AGC	AAC	CAC	TGG	GTG	GTC	AAT	CCG	CGG	CAC	AAC	CTG	777
	Ile	Thr	Ala	Thr	Ser	Asn	His	Trp	Val	Val	Asn	Pro	Arg	His	Asn	Leu	
			230					235					240				
20	GGC	CTG	CAG	CTC	TCG	GTG	GAG	ACG	CTG	GAT	GGG	CAG	AGC	ATC	AAC	CCC	825
	Gly	Leu	Gln	Leu	Ser	Val	Glu	Thr	Leu	Asp	Gly	Gln	Ser	Ile	Asn	Pro	
		245					250					255					
25	AAG	TTG	GCG	GGC	CTG	ATT	GGG	CGG	CAC	GGG	CCC	CAG	AAC	AAG	CAG	CCC	873
	Lys	Leu	Ala	Gly	Leu	Ile	Gly	Arg	His	Gly	Pro	Gln	Asn	Lys	Gln	Pro	
	260					265					270					275	
30	TTC	ATG	GTG	GCT	TTC	TTC	AAG	GCC	ACG	GAG	GTC	CAC	TTC	CGC	AGC	ATC	921
	Phe	Met	Val	Ala	Phe	Phe	Lys	Ala	Thr	Glu	Val	His	Phe	Arg	Ser	Ile	
					280					285					290		
35	CGG	TCC	ACG	GGG	AGC	AAA	CAG	CGC	AGC	CAG	AAC	CGC	TCC	AAG	ACG	CCC	969
	Arg	Ser	Thr	Gly	Ser	Lys	Gln	Arg	Ser	Gln	Asn	Arg	Ser	Lys	Thr	Pro	
				295					300					305			
40	AAG	AAC	CAG	GAA	GCC	CTG	CGG	ATG	GCC	AAC	GTG	GCA	GAG	AAC	AGC	AGC	1017
	Lys	Asn	Gln	Glu	Ala	Leu	Arg	Met	Ala	Asn	Val	Ala	Glu	Asn	Ser	Ser	
			310					315					320				
45	AGC	GAC	CAG	AGG	CAG	GCC	TGT	AAG	AAG	CAC	GAG	CTG	TAT	GTC	AGC	TTC	1065
	Ser	Asp	Gln	Arg	Gln	Ala	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val	Ser	Phe	
		325					330					335					
50	CGA	GAC	CTG	GGC	TGG	CAG	GAC	TGG	ATC	ATC	GCG	CCT	GAA	GGC	TAC	GCC	1113
	Arg	Asp	Leu	Gly	Trp	Gln	Asp	Trp	Ile	Ile	Ala	Pro	Glu	Gly	Tyr	Ala	
		340				345					350					355	
55	GCC	TAC	TAC	TGT	GAG	GGG	GAG	TGT	GCC	TTC	CCT	CTG	AAC	TCC	TAC	ATG	1161
	Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ala	Phe	Pro	Leu	Asn	Ser	Tyr	Met	
					360					365						370	

- 112 -

	AAC GCC ACC AAC CAC GCC ATC GTG CAG ACG CTG GTC CAC TTC ATC AAC	1209
	Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn	
	375 380 385	
5	CCG GAA ACG GTG CCC AAG CCC TGC TGT GCG CCC ACG CAG CTC AAT GCC	1257
	Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala	
	390 395 400	
10	ATC TCC GTC CTC TAC TTC GAT GAC AGC TCC AAC GTC ATC CTG AAG AAA	1305
	Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys	
	405 410 415	
15	TAC AGA AAC ATG GTG GTC CGG GCC TGT GGC TGC CAC TAGCTCCTCC	1351
	Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His	
	420 425 430	
	GAGAATTCAG ACCCTTTGGG GCCAAGTTTT TCTGGATCCT CCATTGCTCG CCTTGGCCAG	1411
20	GAACCAGCAG ACCAACTGCC TTTTGTGAGA CCTTCCCCTC CCTATCCCCA ACTTTAAAGG	1471
	TGTGAGAGTA TTAGGAAACA TGAGCAGCAT ATGGCTTTTG ATCAGTTTTT CAGTGGCAGC	1531
	ATCCAATGAA CAAGATCCTA CAAGCTGTGC AGGCAAAACC TAGCAGGAAA AAAAAACAAC	1591
25	GCATAAAGAA AAATGGCCGG GCCAGGTCAT TGGCTGGGAA GTCTCAGCCA TGCACGGACT	1651
	CGTTTCCAGA GGTAATTATG AGCGCCTACC AGCCAGGCCA CCCAGCCGTG GGAGGAAGGG	1711
30	GGCGTGGCAA GGGGTGGGCA CATTGGTGTC TGTGCGAAAG GAAAATTGAC CCGGAAGTTC	1771
	CTGTAATAAA TGTCACAATA AAACGAATGA ATGAAAAAAAA AAAAAAAAAA A	1822

(2) INFORMATION FOR SEQ ID NO:17:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 431 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

45	Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala
	1 5 10 15
	Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser
	20 25 30

50

- 113 -

Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser
 35 40 45
 5 Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu
 50 55 60
 Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro
 65 70 75 80
 10 Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly
 85 90 95
 Gly Pro Gly Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser
 100 105 110
 15 Thr Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr
 115 120 125
 Asp Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys
 20 130 135 140
 Glu Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu
 145 150 155 160
 25 Ser Lys Ile Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile
 165 170 175
 Tyr Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Arg Ile
 180 185 190
 30 Ser Val Tyr Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu
 195 200 205
 Phe Leu Leu Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu
 35 210 215 220
 Val Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg
 225 230 235 240
 40 His Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser
 245 250 255
 Ile Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn
 260 265 270
 45 Lys Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe
 275 280 285
 Arg Ser Ile Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser
 50 290 295 300

- 114 -

Lys Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu
 305 310 315 320
 Asn Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr
 5 325 330 335
 Val Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu
 340 345 350
 10 Gly Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn
 355 360 365
 Ser Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His
 15 370 375 380
 Phe Ile Asn Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln
 385 390 395 400
 Leu Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile
 20 405 410 415
 Leu Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 420 425 430

25 (2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1873 base pairs
 (B) TYPE: nucleic acid
 30 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

35 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- 40 (A) ORGANISM: MURIDAE
 (F) TISSUE TYPE: EMBRYO

(ix) FEATURE:

- 45 (A) NAME/KEY: CDS
 (B) LOCATION: 104..1393
 (D) OTHER INFORMATION: /function= "OSTEOGENIC PROTEIN"
 /product= "MOP1"
 /note= "MOP1 (CDNA)"

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

- 115 -

	CTGCAGCAAG	TGACCTCGGG	TCGTGGACCG	CTGCCCTGCC	CCCTCCGCTG	CCACCTGGGG	60
	CGGCGCGGGC	CCGGTGCCCC	GGATCGCGCG	TAGAGCCGGC	GCG ATG CAC GTG CGC		115
					Met His Val Arg		
5					1		
	TCG CTG CGC GCT GCG GCG CCA CAC AGC TTC GTG GCG CTC TGG GCG CCT						163
	Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala Leu Trp Ala Pro						
	5 10 15 20						
10	CTG TTC TTG CTG CGC TCC GCC CTG GCC GAT TTC AGC CTG GAC AAC GAG						211
	Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser Leu Asp Asn Glu						
	25 30 35						
15	GTG CAC TCC AGC TTC ATC CAC CGG CGC CTC CGC AGC CAG GAG CGG CGG						259
	Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser Gln Glu Arg Arg						
	40 45 50						
20	GAG ATG CAG CGG GAG ATC CTG TCC ATC TTA GGG TTG CCC CAT CGC CCG						307
	Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu Pro His Arg Pro						
	55 60 65						
25	CGC CCG CAC CTC CAG GGA AAG CAT AAT TCG GCG CCC ATG TTC ATG TTG						355
	Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro Met Phe Met Leu						
	70 75 80						
30	GAC CTG TAC AAC GCC ATG GCG GTG GAG GAG AGC GGG CCG GAC GGA CAG						403
	Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Ser Gly Pro Asp Gly Gln						
	85 90 95 100						
35	GGC TTC TCC TAC CCC TAC AAG GCC GTC TTC AGT ACC CAG GGC CCC CCT						451
	Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser Thr Gln Gly Pro Pro						
	105 110 115						
40	TTA GCC AGC CTG CAG GAC AGC CAT TTC CTC ACT GAC GCC GAC ATG GTC						499
	Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr Asp Ala Asp Met Val						
	120 125 130						
45	ATG AGC TTC GTC AAC CTA GTG GAA CAT GAC AAA GAA TTC TTC CAC CCT						547
	Met Ser Phe Val Asn Leu Val Glu His Asp Lys Glu Phe Phe His Pro						
	135 140 145						
50	CGA TAC CAC CAT CGG GAG TTC CGG TTT GAT CTT TCC AAG ATC CCC GAG						595
	Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu Ser Lys Ile Pro Glu						
	150 155 160						
55	GGC GAA CGG GTG ACC GCA GCC GAA TTC AGG ATC TAT AAG GAC TAC ATC						643
	Gly Glu Arg Val Thr Ala Ala Glu Phe Arg Ile Tyr Lys Asp Tyr Ile						
	165 170 175 180						

- 116 -

	CGG	GAG	CGA	TTT	GAC	AAC	GAG	ACC	TTC	CAG	ATC	ACA	GTC	TAT	CAG	GTG	691
	Arg	Glu	Arg	Phe	Asp	Asn	Glu	Thr	Phe	Gln	Ile	Thr	Val	Tyr	Gln	Val	
					185					190					195		
5	CTC	CAG	GAG	CAC	TCA	GGC	AGG	GAG	TCG	GAC	CTC	TTC	TTG	CTG	GAC	AGC	739
	Leu	Gln	Glu	His	Ser	Gly	Arg	Glu	Ser	Asp	Leu	Phe	Leu	Leu	Asp	Ser	
				200					205					210			
10	CGC	ACC	ATC	TGG	GCT	TCT	GAG	GAG	GGC	TGG	TTG	GTG	TTT	GAT	ATC	ACA	787
	Arg	Thr	Ile	Trp	Ala	Ser	Glu	Glu	Gly	Trp	Leu	Val	Phe	Asp	Ile	Thr	
			215					220					225				
15	GCC	ACC	AGC	AAC	CAC	TGG	GTG	GTC	AAC	CCT	CGG	CAC	AAC	CTG	GGC	TTA	835
	Ala	Thr	Ser	Asn	His	Trp	Val	Val	Asn	Pro	Arg	His	Asn	Leu	Gly	Leu	
		230					235					240					
20	CAG	CTC	TCT	GTG	GAG	ACC	CTG	GAT	GGG	CAG	AGC	ATC	AAC	CCC	AAG	TTG	883
	Gln	Leu	Ser	Val	Glu	Thr	Leu	Asp	Gly	Gln	Ser	Ile	Asn	Pro	Lys	Leu	
	245				250					255						260	
	GCA	GGC	CTG	ATT	GGA	CGG	CAT	GGA	CCC	CAG	AAC	AAG	CAA	CCC	TTC	ATG	931
	Ala	Gly	Leu	Ile	Gly	Arg	His	Gly	Pro	Gln	Asn	Lys	Gln	Pro	Phe	Met	
				265					270						275		
25	GTG	GCC	TTC	TTC	AAG	GCC	ACG	GAA	GTC	CAT	CTC	CGT	AGT	ATC	CGG	TCC	979
	Val	Ala	Phe	Phe	Lys	Ala	Thr	Glu	Val	His	Leu	Arg	Ser	Ile	Arg	Ser	
				280				285						290			
30	ACG	GGG	GGC	AAG	CAG	CGC	AGC	CAG	AAT	CGC	TCC	AAG	ACG	CCA	AAG	AAC	1027
	Thr	Gly	Gly	Lys	Gln	Arg	Ser	Gln	Asn	Arg	Ser	Lys	Thr	Pro	Lys	Asn	
			295					300					305				
35	CAA	GAG	GCC	CTG	AGG	ATG	GCC	AGT	GTG	GCA	GAA	AAC	AGC	AGC	AGT	GAC	1075
	Gln	Glu	Ala	Leu	Arg	Met	Ala	Ser	Val	Ala	Glu	Asn	Ser	Ser	Ser	Asp	
		310					315					320					
40	CAG	AGG	CAG	GCC	TGC	AAG	AAA	CAT	GAG	CTG	TAC	GTC	AGC	TTC	CGA	GAC	1123
	Gln	Arg	Gln	Ala	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val	Ser	Phe	Arg	Asp	
	325				330					335						340	
	CTT	GGC	TGG	CAG	GAC	TGG	ATC	ATT	GCA	CCT	GAA	GGC	TAT	GCT	GCC	TAC	1171
	Leu	Gly	Trp	Gln	Asp	Trp	Ile	Ile	Ala	Pro	Glu	Gly	Tyr	Ala	Ala	Tyr	
				345					350					355			
45	TAC	TGT	GAG	GGA	GAG	TGC	GCC	TTC	CCT	CTG	AAC	TCC	TAC	ATG	AAC	GCC	1219
	Tyr	Cys	Glu	Gly	Glu	Cys	Ala	Phe	Pro	Leu	Asn	Ser	Tyr	Met	Asn	Ala	
				360					365					370			

- 117 -

ACC AAC CAC GCC ATC GTC CAG ACA CTG GTT CAC TTC ATC AAC CCA GAC 1267
 Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn Pro Asp
 375 380 385

5 ACA GTA CCC AAG CCC TGC TGT GCG CCC ACC CAG CTC AAC GCC ATC TCT 1315
 Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala Ile Ser
 390 395 400

10 GTC CTC TAC TTC GAC GAC AGC TCT AAT GTC GAC CTG AAG AAG TAC AGA 1363
 Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Asp Leu Lys Lys Tyr Arg
 405 410 415 420

15 AAC ATG GTG GTC CGG GCC TGT GGC TGC CAC TAGCTCTTCC TGAGACCCTG 1413
 Asn Met Val Val Arg Ala Cys Gly Cys His
 425 430

ACCTTTGCGG GGCCACACCT TTCCAAATCT TCGATGTCTC ACCATCTAAG TCTCTCACTG 1473

20 CCCACCTTGG CGAGGAGAAC AGACCAACCT CTCCTGAGCC TTCCCTCACC TCCCAACCGG 1533

AAGCATGTAA GGGTTCAGG AACCTGAGCG TGCAGCAGCT GATGAGCGCC CTTTCCTTCT 1593

GGCACGTGAC GGACAAGATC CTACCAGCTA CCACAGCAAA CGCCTAAGAG CAGGAAAAAT 1653

25 GTCTGCCAGG AAAGTGTCCA GTGTCCACAT GGCCCCTGGC GCTCTGAGTC TTTGAGGAGT 1713

AATCGCAAGC CTCGTTTACG TGCAGCAGAA GGAAGGGCTT AGCCAGGGTG GCGGCTGGCG 1773

TCTGTGTTGA AGGGAAACCA AGCAGAAGCC ACTGTAATGA TATGTCACAA TAAAACCCAT 1833

30 GAATGAAAAA AAAAAAAAAA AAAAAAAAAA AAAAGAATTTC 1873

(2) INFORMATION FOR SEQ ID NO:19:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 430 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

45 Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala
 1 5 10 15

Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser
 20 25 30

50

- 118 -

Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser
 35 40 45
 5 Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu
 50 55 60
 Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro
 65 70 75 80
 10 Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Ser Gly
 85 90 95
 Pro Asp Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser Thr
 100 105 110
 15 Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr Asp
 115 120 125
 Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys Glu
 20 130 135 140
 Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu Ser
 145 150 155 160
 25 Lys Ile Pro Glu Gly Glu Arg Val Thr Ala Ala Glu Phe Arg Ile Tyr
 165 170 175
 Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Gln Ile Thr
 180 185 190
 30 Val Tyr Gln Val Leu Gln Glu His Ser Gly Arg Glu Ser Asp Leu Phe
 195 200 205
 Leu Leu Asp Ser Arg Thr Ile Trp Ala Ser Glu Glu Gly Trp Leu Val
 35 210 215 220
 Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg His
 225 230 235 240
 40 Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser Ile
 245 250 255
 Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn Lys
 260 265 270
 45 Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Leu Arg
 275 280 285

- 119 -

Ser Ile Arg Ser Thr Gly Gly Lys Gln Arg Ser Gln Asn Arg Ser Lys
 290 295 300
 Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Ser Val Ala Glu Asn
 5 305 310 315 320
 Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val
 325 330 335
 10 Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly
 340 345 350
 Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser
 355 360 365
 15 Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe
 370 375 380
 20 Ile Asn Pro Asp Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu
 385 390 395 400
 Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Asp Leu
 405 410 415
 25 Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 420 425 430

(2) INFORMATION FOR SEQ ID NO:20:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1723 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 35
 (ii) MOLECULE TYPE: cDNA
 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens
 40 (F) TISSUE TYPE: HIPPOCAMPUS
 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 490..1696
 45 (D) OTHER INFORMATION: /function= "OSTEOGENIC PROTEIN"
 /product= "hOP2-PP"
 /note= "hOP2 (cDNA)"
 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

- 120 -

	GGCGCCGGCA GAGCAGGAGT GGCTGGAGGA GCTGTGGTTG GAGCAGGAGG TGGCACGGCA	60
	GGGCTGGAGG GCTCCCTATG AGTGGCGGAG ACGGCCAGG AGGCGCTGGA GCAACAGCTC	120
5	CCACACCGCA CCAAGCGGTG GCTGCAGGAG CTCGCCCATC GCCCCTGCGC TGCTCGGACC	180
	GCGGCCACAG CCGGACTGGC GGGTACGGCG GCGACAGAGG CATTGGCCGA GAGTCCCAGT	240
	CCGCAGAGTA GCCCCGGCCT CGAGGCGGTG GCGTCCCGGT CCTCTCCGTC CAGGAGCCAG	300
10	GACAGGTGTC GCGCGGCGGG GCTCCAGGGA CCGCGCCTGA GGCCGGCTGC CCGCCCGTCC	360
	CGCCCCGCCC CGCCGCCCCG CGCCGCCCCA GCCCAGCCTC CTTGCCGTCG GGGCGTCCCC	420
15	AGGCCCTGGG TCGGCCGCGG AGCCGATGCG CGCCGCTGA GCGCCCCAGC TGAGCGCCCC	480
	CGGCCTGCC ATG ACC GCG CTC CCC GGC CCG CTC TGG CTC CTG GGC CTG	528
	Met Thr Ala Leu Pro Gly Pro Leu Trp Leu Leu Gly Leu	
	1 5 10	
20	GCG CTA TGC GCG CTG GGC GGG GGC GGC CCC GGC CTG CGA CCC CCG CCC	576
	Ala Leu Cys Ala Leu Gly Gly Gly Gly Pro Gly Leu Arg Pro Pro Pro	
	15 20 25	
25	GGC TGT CCC CAG CGA CGT CTG GGC GCG CGC GAG CGC CGG GAC GTG CAG	624
	Gly Cys Pro Gln Arg Arg Leu Gly Ala Arg Glu Arg Arg Asp Val Gln	
	30 35 40 45	
30	CGC GAG ATC CTG GCG GTG CTC GGG CTG CCT GGG CGG CCC CGG CCC CGC	672
	Arg Glu Ile Leu Ala Val Leu Gly Leu Pro Gly Arg Pro Arg Pro Arg	
	50 55 60	
35	GCG CCA CCC GCC GCC TCC CGG CTG CCC GCG TCC GCG CCG CTC TTC ATG	720
	Ala Pro Pro Ala Ala Ser Arg Leu Pro Ala Ser Ala Pro Leu Phe Met	
	65 70 75	
	CTG GAC CTG TAC CAC GCC ATG GCC GGC GAC GAC GAC GAG GAC GGC GCG	768
	Leu Asp Leu Tyr His Ala Met Ala Gly Asp Asp Asp Glu Asp Gly Ala	
	80 85 90	
40	CCC GCG GAG CGG CGC CTG GGC CGC GCC GAC CTG GTC ATG AGC TTC GTT	816
	Pro Ala Glu Arg Arg Leu Gly Arg Ala Asp Leu Val Met Ser Phe Val	
	95 100 105	
45	AAC ATG GTG GAG CGA GAC CGT GCC CTG GGC CAC CAG GAG CCC CAT TGG	864
	Asn Met Val Glu Arg Asp Arg Ala Leu Gly His Gln Glu Pro His Trp	
	110 115 120 125	

- 121 -

	AAG GAG TTC CGC TTT GAC CTG ACC CAG ATC CCG GCT GGG GAG GCG GTC	912
	Lys Glu Phe Arg Phe Asp Leu Thr Gln Ile Pro Ala Gly Glu Ala Val	
	130 135 140	
5	ACA GCT GCG GAG TTC CGG ATT TAC AAG GTG CCC AGC ATC CAC CTG CTC	960
	Thr Ala Ala Glu Phe Arg Ile Tyr Lys Val Pro Ser Ile His Leu Leu	
	145 150 155	
10	AAC AGG ACC CTC CAC GTC AGC ATG TTC CAG GTG GTC CAG GAG CAG TCC	1008
	Asn Arg Thr Leu His Val Ser Met Phe Gln Val Val Gln Glu Gln Ser	
	160 165 170	
15	AAC AGG GAG TCT GAC TTG TTC TTT TTG GAT CTT CAG ACG CTC CGA GCT	1056
	Asn Arg Glu Ser Asp Leu Phe Phe Leu Asp Leu Gln Thr Leu Arg Ala	
	175 180 185	
20	GGA GAC GAG GGC TGG CTG GTG CTG GAT GTC ACA GCA GCC AGT GAC TGC	1104
	Gly Asp Glu Gly Trp Leu Val Leu Asp Val Thr Ala Ala Ser Asp Cys	
	190 195 200 205	
20	TGG TTG CTG AAG CGT CAC AAG GAC CTG GGA CTC CGC CTC TAT GTG GAG	1152
	Trp Leu Leu Lys Arg His Lys Asp Leu Gly Leu Arg Leu Tyr Val Glu	
	210 215 220	
25	ACT GAG GAC GGG CAC AGC GTG GAT CCT GGC CTG GCC GGC CTG CTG GGT	1200
	Thr Glu Asp Gly His Ser Val Asp Pro Gly Leu Ala Gly Leu Leu Gly	
	225 230 235	
30	CAA CGG GCC CCA CGC TCC CAA CAG CCT TTC GTG GTC ACT TTC TTC AGG	1248
	Gln Arg Ala Pro Arg Ser Gln Gln Pro Phe Val Val Thr Phe Phe Arg	
	240 245 250	
35	GCC AGT CCG AGT CCC ATC CGC ACC CCT CGG GCA GTG AGG CCA CTG AGG	1296
	Ala Ser Pro Ser Pro Ile Arg Thr Pro Arg Ala Val Arg Pro Leu Arg	
	255 260 265	
40	AGG AGG CAG CCG AAG AAA AGC AAC GAG CTG CCG CAG GCC AAC CGA CTC	1344
	Arg Arg Gln Pro Lys Lys Ser Asn Glu Leu Pro Gln Ala Asn Arg Leu	
	270 275 280 285	
40	CCA GGG ATC TTT GAT GAC GTC CAC GGC TCC CAC GGC CGG CAG GTC TGC	1392
	Pro Gly Ile Phe Asp Asp Val His Gly Ser His Gly Arg Gln Val Cys	
	290 295 300	
45	CGT CGG CAC GAG CTC TAC GTC AGC TTC CAG GAC CTC GGC TGG CTG GAC	1440
	Arg Arg His Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Leu Asp	
	305 310 315	

- 122 -

TGG GTC ATC GCT CCC CAA GGC TAC TCG GCC TAT TAC TGT GAG GGG GAG 1488
 Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala Tyr Tyr Cys Glu Gly Glu
 320 325 330
 5 TGC TCC TTC CCA CTG GAC TCC TGC ATG AAT GCC ACC AAC CAC GCC ATC 1536
 Cys Ser Phe Pro Leu Asp Ser Cys Met Asn Ala Thr Asn His Ala Ile
 335 340 345
 10 CTG CAG TCC CTG GTG CAC CTG ATG AAG CCA AAC GCA GTC CCC AAG GCG 1584
 Leu Gln Ser Leu Val His Leu Met Lys Pro Asn Ala Val Pro Lys Ala
 350 355 360 365
 15 TGC TGT GCA CCC ACC AAG CTG AGC GCC ACC TCT GTG CTC TAC TAT GAC 1632
 Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr Ser Val Leu Tyr Tyr Asp
 370 375 380
 AGC AGC AAC AAC GTC ATC CTG CGC AAA GCC CGC AAC ATG GTG GTC AAG 1680
 Ser Ser Asn Asn Val Ile Leu Arg Lys Ala Arg Asn Met Val Val Lys
 385 390 395
 20 GCC TGC GGC TGC CAC T GAGTCAGCCC GCCCAGCCCT ACTGCAG 1723
 Ala Cys Gly Cys His
 400

25

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 402 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Met Thr Ala Leu Pro Gly Pro Leu Trp Leu Leu Gly Leu Ala Leu Cys
 1 5 10 15
 40 Ala Leu Gly Gly Gly Gly Pro Gly Leu Arg Pro Pro Pro Gly Cys Pro
 20 25 30
 Gln Arg Arg Leu Gly Ala Arg Glu Arg Arg Asp Val Gln Arg Glu Ile
 35 40 45
 45 Leu Ala Val Leu Gly Leu Pro Gly Arg Pro Arg Pro Arg Ala Pro Pro
 50 55 60

- 123 -

Ala Ala Ser Arg Leu Pro Ala Ser Ala Pro Leu Phe Met Leu Asp Leu
 65 70 75 80
 Tyr His Ala Met Ala Gly Asp Asp Asp Glu Asp Gly Ala Pro Ala Glu
 5 85 90 95
 Arg Arg Leu Gly Arg Ala Asp Leu Val Met Ser Phe Val Asn Met Val
 100 105 110
 10 Glu Arg Asp Arg Ala Leu Gly His Gln Glu Pro His Trp Lys Glu Phe
 115 120 125
 Arg Phe Asp Leu Thr Gln Ile Pro Ala Gly Glu Ala Val Thr Ala Ala
 130 135 140
 15 Glu Phe Arg Ile Tyr Lys Val Pro Ser Ile His Leu Leu Asn Arg Thr
 145 150 155 160
 Leu His Val Ser Met Phe Gln Val Val Gln Glu Gln Ser Asn Arg Glu
 165 170 175
 20 Ser Asp Leu Phe Phe Leu Asp Leu Gln Thr Leu Arg Ala Gly Asp Glu
 180 185 190
 25 Gly Trp Leu Val Leu Asp Val Thr Ala Ala Ser Asp Cys Trp Leu Leu
 195 200 205
 Lys Arg His Lys Asp Leu Gly Leu Arg Leu Tyr Val Glu Thr Glu Asp
 210 215 220
 30 Gly His Ser Val Asp Pro Gly Leu Ala Gly Leu Leu Gly Gln Arg Ala
 225 230 235 240
 Pro Arg Ser Gln Gln Pro Phe Val Val Thr Phe Phe Arg Ala Ser Pro
 245 250 255
 35 Ser Pro Ile Arg Thr Pro Arg Ala Val Arg Pro Leu Arg Arg Arg Gln
 260 265 270
 40 Pro Lys Lys Ser Asn Glu Leu Pro Gln Ala Asn Arg Leu Pro Gly Ile
 275 280 285
 Phe Asp Asp Val His Gly Ser His Gly Arg Gln Val Cys Arg Arg His
 290 295 300
 45 Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Leu Asp Trp Val Ile
 305 310 315 320
 Ala Pro Gln Gly Tyr Ser Ala Tyr Tyr Cys Glu Gly Glu Cys Ser Phe
 325 330 335
 50

Pro Leu Asp Ser Cys Met Asn Ala Thr Asn His Ala Ile Leu Gln Ser
340 345 350

Leu Val His Leu Met Lys Pro Asn Ala Val Pro Lys Ala Cys Cys Ala
5 355 360 365

Pro Thr Lys Leu Ser Ala Thr Ser Val Leu Tyr Tyr Asp Ser Ser Asn
370 375 380

10 Asn Val Ile Leu Arg Lys Ala Arg Asn Met Val Val Lys Ala Cys Gly
385 390 395 400

Cys His

15

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 1926 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vi) ORIGINAL SOURCE:

25 (A) ORGANISM: MURIDAE
(F) TISSUE TYPE: EMBRYO

(ix) FEATURE:

30 (A) NAME/KEY: CDS
(B) LOCATION: 93..1289
(D) OTHER INFORMATION: /function= "OSTEOGENIC PROTEIN"
/product= "mOP2-PP"
/note= "mOP2 cDNA"

35

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CCCAGGCACA GGTGCCCGT CTGGTCCTCC CCGTCTGGCG TCAGCCGAGC CCGACCAGCT 60

40 ACCAGTGGAT GCGCGCCGGC TGAAAGTCCG AG ATG GCT ATG CGT CCC GGG CCA 113
Met Ala Met Arg Pro Gly Pro
1 5

45 CTC TGG CTA TTG GGC CTT GCT CTG TGC GCG CTG GGA GGC GGC CAC GGT 161
Leu Trp Leu Leu Gly Leu Ala Leu Cys Ala Leu Gly Gly Gly His Gly
10 15 20

50 CCG CGT CCC CCG CAC ACC TGT CCC CAG CGT CGC CTG GGA GCG CGC GAG 209
Pro Arg Pro Pro His Thr Cys Pro Gln Arg Arg Leu Gly Ala Arg Glu
25 30 35

- 125 -

	CGC	CGC	GAC	ATG	CAG	CGT	GAA	ATC	CTG	GCG	GTG	CTC	GGG	CTA	CCG	GGA	257
	Arg	Arg	Asp	Met	Gln	Arg	Glu	Ile	Leu	Ala	Val	Leu	Gly	Leu	Pro	Gly	
	40					45					50					55	
5	CGG	CCC	CGA	CCC	CGT	GCA	CAA	CCC	GCC	GCT	GCC	CGG	CAG	CCA	GCG	TCC	305
	Arg	Pro	Arg	Pro	Arg	Ala	Gln	Pro	Ala	Ala	Ala	Arg	Gln	Pro	Ala	Ser	
					60					65					70		
10	GCG	CCC	CTC	TTC	ATG	TTG	GAC	CTA	TAC	CAC	GCC	ATG	ACC	GAT	GAC	GAC	353
	Ala	Pro	Leu	Phe	Met	Leu	Asp	Leu	Tyr	His	Ala	Met	Thr	Asp	Asp	Asp	
				75					80					85			
15	GAC	GGC	GGG	CCA	CCA	CAG	GCT	CAC	TTA	GGC	CGT	GCC	GAC	CTG	GTC	ATG	401
	Asp	Gly	Gly	Pro	Pro	Gln	Ala	His	Leu	Gly	Arg	Ala	Asp	Leu	Val	Met	
			90					95					100				
20	AGC	TTC	GTC	AAC	ATG	GTG	GAA	CGC	GAC	CGT	ACC	CTG	GGC	TAC	CAG	GAG	449
	Ser	Phe	Val	Asn	Met	Val	Glu	Arg	Asp	Arg	Thr	Leu	Gly	Tyr	Gln	Glu	
		105					110					115					
25	CCA	CAC	TGG	AAG	GAA	TTC	CAC	TTT	GAC	CTA	ACC	CAG	ATC	CCT	GCT	GGG	497
	Pro	His	Trp	Lys	Glu	Phe	His	Phe	Asp	Leu	Thr	Gln	Ile	Pro	Ala	Gly	
		120				125					130					135	
30	GAG	GCT	GTC	ACA	GCT	GCT	GAG	TTC	CGG	ATC	TAC	AAA	GAA	CCC	AGC	ACC	545
	Glu	Ala	Val	Thr	Ala	Ala	Glu	Phe	Arg	Ile	Tyr	Lys	Glu	Pro	Ser	Thr	
					140					145					150		
35	CAC	CCG	CTC	AAC	ACA	ACC	CTC	CAC	ATC	AGC	ATG	TTC	GAA	GTG	GTC	CAA	593
	His	Pro	Leu	Asn	Thr	Thr	Leu	His	Ile	Ser	Met	Phe	Glu	Val	Val	Gln	
				155					160					165			
40	GAG	CAC	TCC	AAC	AGG	GAG	TCT	GAC	TTG	TTC	TTT	TTG	GAT	CTT	CAG	ACG	641
	Glu	His	Ser	Asn	Arg	Glu	Ser	Asp	Leu	Phe	Phe	Leu	Asp	Leu	Gln	Thr	
			170					175					180				
45	CTC	CGA	TCT	GGG	GAC	GAG	GGC	TGG	CTG	GTG	CTG	GAC	ATC	ACA	GCA	GCC	689
	Leu	Arg	Ser	Gly	Asp	Glu	Gly	Trp	Leu	Val	Leu	Asp	Ile	Thr	Ala	Ala	
		185					190					195					
50	AGT	GAC	CGA	TGG	CTG	CTG	AAC	CAT	CAC	AAG	GAC	CTG	GGA	CTC	CGC	CTC	737
	Ser	Asp	Arg	Trp	Leu	Leu	Asn	His	His	Lys	Asp	Leu	Gly	Leu	Arg	Leu	
		200				205					210					215	
55	TAT	GTG	GAA	ACC	GCG	GAT	GGG	CAC	AGC	ATG	GAT	CCT	GGC	CTG	GCT	GGT	785
	Tyr	Val	Glu	Thr	Ala	Asp	Gly	His	Ser	Met	Asp	Pro	Gly	Leu	Ala	Gly	
					220					225					230		

- 126 -

	CTG CTT GGA CGA CAA GCA CCA CGC TCC AGA CAG CCT TTC ATG GTA ACC	833
	Leu Leu Gly Arg Gln Ala Pro Arg Ser Arg Gln Pro Phe Met Val Thr	
	235 240 245	
5	TTC TTC AGG GCC AGC CAG AGT CCT GTG CGG GCC CCT CGG GCA GCG AGA	881
	Phe Phe Arg Ala Ser Gln Ser Pro Val Arg Ala Pro Arg Ala Ala Arg	
	250 255 260	
10	CCA CTG AAG AGG AGG CAG CCA AAG AAA ACG AAC GAG CTT CCG CAC CCC	929
	Pro Leu Lys Arg Arg Gln Pro Lys Lys Thr Asn Glu Leu Pro His Pro	
	265 270 275	
15	AAC AAA CTC CCA GGG ATC TTT GAT GAT GGC CAC GGT TCC CGC GGC AGA	977
	Asn Lys Leu Pro Gly Ile Phe Asp Asp Gly His Gly Ser Arg Gly Arg	
	280 285 290 295	
20	GAG GTT TGC CGC AGG CAT GAG CTC TAC GTC AGC TTC CGT GAC CTT GGC	1025
	Glu Val Cys Arg Arg His Glu Leu Tyr Val Ser Phe Arg Asp Leu Gly	
	300 305 310	
25	TGG CTG GAC TGG GTC ATC GCC CCC CAG GGC TAC TCT GCC TAT TAC TGT	1073
	Trp Leu Asp Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala Tyr Tyr Cys	
	315 320 325	
30	GAG GGG GAG TGT GCT TTC CCA CTG GAC TCC TGT ATG AAC GCC ACC AAC	1121
	Glu Gly Glu Cys Ala Phe Pro Leu Asp Ser Cys Met Asn Ala Thr Asn	
	330 335 340	
35	CAT GCC ATC TTG CAG TCT CTG GTG CAC CTG ATG AAG CCA GAT GTT GTC	1169
	His Ala Ile Leu Gln Ser Leu Val His Leu Met Lys Pro Asp Val Val	
	345 350 355	
40	CCC AAG GCA TGC TGT GCA CCC ACC AAA CTG AGT GCC ACC TCT GTG CTG	1217
	Pro Lys Ala Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr Ser Val Leu	
	360 365 370 375	
45	TAC TAT GAC AGC AGC AAC AAT GTC ATC CTG CGT AAA CAC CGT AAC ATG	1265
	Tyr Tyr Asp Ser Ser Asn Asn Val Ile Leu Arg Lys His Arg Asn Met	
	380 385 390	
50	GTG GTC AAG GCC TGT GGC TGC CAC TGAGGCCCCG CCCAGCATCC TGCTTCTACT	1319
	Val Val Lys Ala Cys Gly Cys His	
	395	
45	ACCTTACCAT CTGGCCGGGC CCCTCTCCAG AGGCAGAAAC CCTTCTATGT TATCATAGCT	1379
	CAGACAGGGG CAATGGGAGG CCCTTCACTT CCCCTGGCCA CTTCCTGCTA AAATTCTGGT	1439
	CTTTCCAGT TCCTCTGTCC TTCATGGGGT TTCGGGGCTA TCACCCCGCC CTCTCCATCC	1499

- 127 -

TCCTACCCCA AGCATAGACT GAATGCACAC AGCATCCCAG AGCTATGCTA ACTGAGAGGT 1559
 CTGGGGTCAG CACTGAAGGC CCACATGAGG AAGACTGATC CTTGGCCATC CTCAGCCCAC 1619
 5 AATGGCAAAT TCTGGATGGT CTAAGAAGGC CCTGGAATTC TAAACTAGAT GATCTGGGCT 1679
 CTCTGCACCA TTCATTGTGG CAGTTGGGAC ATTTTITAGGT ATAACAGACA CATACTTA 1739
 GATCAATGCA TCGCTGTACT CCTTGAAATC AGAGCTAGCT TGTTAGAAAA AGAATCAGAG 1799
 10 CCAGGTATAG CGGTGCATGT CATTAAATCC AGCGCTAAAG AGACAGAGAC AGGAGAATCT 1859
 CTGTGAGTTC AAGGCCACAT AGAAAGAGCC TGTCTCGGGA GCAGGAAAAA AAAAAAAAC 1919
 15 GGAATTC 1926

(2) INFORMATION FOR SEQ ID NO:23:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 399 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Met Ala Met Arg Pro Gly Pro Leu Trp Leu Leu Gly Leu Ala Leu Cys
 30 1 5 10 15
 Ala Leu Gly Gly Gly His Gly Pro Arg Pro Pro His Thr Cys Pro Gln
 20 25 30
 35 Arg Arg Leu Gly Ala Arg Glu Arg Arg Asp Met Gln Arg Glu Ile Leu
 35 40 45
 Ala Val Leu Gly Leu Pro Gly Arg Pro Arg Pro Arg Ala Gln Pro Ala
 50 55 60
 40 Ala Ala Arg Gln Pro Ala Ser Ala Pro Leu Phe Met Leu Asp Leu Tyr
 65 70 75 80
 His Ala Met Thr Asp Asp Asp Asp Gly Gly Pro Pro Gln Ala His Leu
 45 85 90 95
 Gly Arg Ala Asp Leu Val Met Ser Phe Val Asn Met Val Glu Arg Asp
 100 105 110

- 128 -

Arg Thr Leu Gly Tyr Gln Glu Pro His Trp Lys Glu Phe His Phe Asp
 115 120 125
 5 Leu Thr Gln Ile Pro Ala Gly Glu Ala Val Thr Ala Ala Glu Phe Arg
 130 135 140
 Ile Tyr Lys Glu Pro Ser Thr His Pro Leu Asn Thr Thr Leu His Ile
 145 150 155 160
 10 Ser Met Phe Glu Val Val Gln Glu His Ser Asn Arg Glu Ser Asp Leu
 165 170 175
 Phe Phe Leu Asp Leu Gln Thr Leu Arg Ser Gly Asp Glu Gly Trp Leu
 180 185 190
 15 Val Leu Asp Ile Thr Ala Ala Ser Asp Arg Trp Leu Leu Asn His His
 195 200 205
 Lys Asp Leu Gly Leu Arg Leu Tyr Val Glu Thr Ala Asp Gly His Ser
 210 215 220
 20 Met Asp Pro Gly Leu Ala Gly Leu Leu Gly Arg Gln Ala Pro Arg Ser
 225 230 235 240
 25 Arg Gln Pro Phe Met Val Thr Phe Phe Arg Ala Ser Gln Ser Pro Val
 245 250 255
 Arg Ala Pro Arg Ala Ala Arg Pro Leu Lys Arg Arg Gln Pro Lys Lys
 260 265 270
 30 Thr Asn Glu Leu Pro His Pro Asn Lys Leu Pro Gly Ile Phe Asp Asp
 275 280 285
 Gly His Gly Ser Arg Gly Arg Glu Val Cys Arg Arg His Glu Leu Tyr
 290 295 300
 35 Val Ser Phe Arg Asp Leu Gly Trp Leu Asp Trp Val Ile Ala Pro Gln
 305 310 315 320
 40 Gly Tyr Ser Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asp
 325 330 335
 Ser Cys Met Asn Ala Thr Asn His Ala Ile Leu Gln Ser Leu Val His
 340 345 350
 45

- 129 -

Leu Met Lys Pro Asp Val Val Pro Lys Ala Cys Cys Ala Pro Thr Lys
 355 360 365

5 Leu Ser Ala Thr Ser Val Leu Tyr Tyr Asp Ser Ser Asn Asn Val Ile
 370 375 380

Leu Arg Lys His Arg Asn Met Val Val Lys Ala Cys Gly Cys His
 385 390 395

10 (2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 1368 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..1368

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

30	1	ATG TCG GGA CTG CGA AAC ACC TCG GAG GCC GTT GCA GTG CTC GCC TCC	48
		Met Ser Gly Leu Arg Asn Thr Ser Glu Ala Val Ala Val Leu Ala Ser	
		5 10 15	
		CTG GGA CTC GGA ATG GTT CTG CTC ATG TTC GTG GCG ACC ACG CCG CCG	96
		Leu Gly Leu Gly Met Val Leu Leu Met Phe Val Ala Thr Thr Pro Pro	
		20 25 30	
35		GCC GTT GAG GCC ACC CAG TCG GGG ATT TAC ATA GAC AAC GGC AAG GAC	144
		Ala Val Glu Ala Thr Gln Ser Gly Ile Tyr Ile Asp Asn Gly Lys Asp	
		35 40 45	
40		CAG ACG ATC ATG CAC AGA GTG CTG AGC GAG GAC GAC AAG CTG GAC GTC	192
		Gln Thr Ile Met His Arg Val Leu Ser Glu Asp Asp Lys Leu Asp Val	
		50 55 60	
45		TCG TAC GAG ATC CTC GAG TTC CTG GGC ATC GCC GAA CGG CCG ACG CAC	240
		Ser Tyr Glu Ile Leu Glu Phe Leu Gly Ile Ala Glu Arg Pro Thr His	
		65 70 75 80	

- 130 -

	CTG	AGC	AGC	CAC	CAG	TTG	TCG	CTG	AGG	AAG	TCG	GCT	CCC	AAG	TTC	CTG	288
	Leu	Ser	Ser	His	Gln	Leu	Ser	Leu	Arg	Lys	Ser	Ala	Pro	Lys	Phe	Leu	
					85					90					95		
5	CTG	GAC	GTC	TAC	CAC	CGC	ATC	ACG	GCG	GAG	GAG	GGT	CTC	AGC	GAT	CAG	336
	Leu	Asp	Val	Tyr	His	Arg	Ile	Thr	Ala	Glu	Glu	Gly	Leu	Ser	Asp	Gln	
				100					105					110			
10	GAT	GAG	GAC	GAC	GAC	TAC	GAA	CGC	GGC	CAT	CGG	TCC	AGG	AGG	AGC	GCC	384
	Asp	Glu	Asp	Asp	Asp	Tyr	Glu	Arg	Gly	His	Arg	Ser	Arg	Arg	Ser	Ala	
				115				120					125				
15	GAC	CTC	GAG	GAG	GAT	GAG	GGC	GAG	CAG	CAG	AAG	AAC	TTC	ATC	ACC	GAC	432
	Asp	Leu	Glu	Glu	Asp	Glu	Gly	Glu	Gln	Gln	Lys	Asn	Phe	Ile	Thr	Asp	
				130			135					140					
20	CTG	GAC	AAG	CGG	GCC	ATC	GAC	GAG	AGC	GAC	ATC	ATC	ATG	ACC	TTC	CTG	480
	Leu	Asp	Lys	Arg	Ala	Ile	Asp	Glu	Ser	Asp	Ile	Ile	Met	Thr	Phe	Leu	
						150					155					160	
25	AAC	AAG	CGC	CAC	CAC	AAT	GTG	GAC	GAA	CTG	CGT	CAC	GAG	CAC	GGC	CGT	528
	Asn	Lys	Arg	His	His	Asn	Val	Asp	Glu	Leu	Arg	His	Glu	His	Gly	Arg	
					165					170					175		
30	CGC	CTG	TGG	TTC	GAC	GTC	TCC	AAC	GTG	CCC	AAC	GAC	AAC	TAC	CTG	GTG	576
	Arg	Leu	Trp	Phe	Asp	Val	Ser	Asn	Val	Pro	Asn	Asp	Asn	Tyr	Leu	Val	
					180				185					190			
35	ATG	GCC	GAG	CTG	CGC	ATC	TAT	CAG	AAC	GCC	AAC	GAG	GGC	AAG	TGG	CTG	624
	Met	Ala	Glu	Leu	Arg	Ile	Tyr	Gln	Asn	Ala	Asn	Glu	Gly	Lys	Trp	Leu	
				195				200					205				
40	ACC	GCC	AAC	AGG	GAG	TTC	ACC	ATC	ACG	GTA	TAC	GCC	ATT	GGC	ACC	GGC	672
	Thr	Ala	Asn	Arg	Glu	Phe	Thr	Ile	Thr	Val	Tyr	Ala	Ile	Gly	Thr	Gly	
				210			215					220					
45	ACG	CTG	GGC	CAG	CAC	ACC	ATG	GAG	CCG	CTG	TCC	TCG	GTG	AAC	ACC	ACC	720
	Thr	Leu	Gly	Gln	His	Thr	Met	Glu	Pro	Leu	Ser	Ser	Val	Asn	Thr	Thr	
						230				235					240		
50	GGG	GAC	TAC	GTG	GGC	TGG	TTG	GAG	CTC	AAC	GTG	ACC	GAG	GGC	CTG	CAC	768
	Gly	Asp	Tyr	Val	Gly	Trp	Leu	Glu	Leu	Asn	Val	Thr	Glu	Gly	Leu	His	
					245					250					255		
55	GAG	TGG	CTG	GTC	AAG	TCG	AAG	GAC	AAT	CAT	GGC	ATC	TAC	ATT	GGA	GCA	816
	Glu	Trp	Leu	Val	Lys	Ser	Lys	Asp	Asn	His	Gly	Ile	Tyr	Ile	Gly	Ala	
					260				265					270			

[illegible]

- 132 -

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 455 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Met Ser Gly Leu Arg Asn Thr Ser Glu Ala Val Ala Val Leu Ala Ser
1 5 10 15

15 Leu Gly Leu Gly Met Val Leu Leu Met Phe Val Ala Thr Thr Pro Pro
20 25 30

Ala Val Glu Ala Thr Gln Ser Gly Ile Tyr Ile Asp Asn Gly Lys Asp
35 40 45

20 Gln Thr Ile Met His Arg Val Leu Ser Glu Asp Asp Lys Leu Asp Val
50 55 60

25 Ser Tyr Glu Ile Leu Glu Phe Leu Gly Ile Ala Glu Arg Pro Thr His
65 70 75 80

Leu Ser Ser His Gln Leu Ser Leu Arg Lys Ser Ala Pro Lys Phe Leu
85 90 95

30 Leu Asp Val Tyr His Arg Ile Thr Ala Glu Glu Gly Leu Ser Asp Gln
100 105 110

Asp Glu Asp Asp Asp Tyr Glu Arg Gly His Arg Ser Arg Arg Ser Ala
115 120 125

35 Asp Leu Glu Glu Asp Glu Gly Glu Gln Gln Lys Asn Phe Ile Thr Asp
130 135 140

40 Leu Asp Lys Arg Ala Ile Asp Glu Ser Asp Ile Ile Met Thr Phe Leu
145 150 155 160

Asn Lys Arg His His Asn Val Asp Glu Leu Arg His Glu His Gly Arg
165 170 175

45 Arg Leu Trp Phe Asp Val Ser Asn Val Pro Asn Asp Asn Tyr Leu Val
180 185 190

Met Ala Glu Leu Arg Ile Tyr Gln Asn Ala Asn Glu Gly Lys Trp Leu
195 200 205

50

- 133 -

Thr Ala Asn Arg Glu Phe Thr Ile Thr Val Tyr Ala Ile Gly Thr Gly
 210 215 220
 5 Thr Leu Gly Gln His Thr Met Glu Pro Leu Ser Ser Val Asn Thr Thr
 225 230 235 240
 Gly Asp Tyr Val Gly Trp Leu Glu Leu Asn Val Thr Glu Gly Leu His
 245 250 255
 10 Glu Trp Leu Val Lys Ser Lys Asp Asn His Gly Ile Tyr Ile Gly Ala
 260 265 270
 His Ala Val Asn Arg Pro Asp Arg Glu Val Lys Leu Asp Asp Ile Gly
 275 280 285
 15 Leu Ile His Arg Lys Val Asp Asp Glu Phe Gln Pro Phe Met Ile Gly
 290 295 300
 20 Phe Phe Arg Gly Pro Glu Leu Ile Lys Ala Thr Ala His Ser Ser His
 305 310 315 320
 His Arg Ser Lys Arg Ser Ala Ser His Pro Arg Lys Arg Lys Lys Ser
 325 330 335
 25 Val Ser Pro Asn Asn Val Pro Leu Leu Glu Pro Met Glu Ser Thr Arg
 340 345 350
 Ser Cys Gln Met Gln Thr Leu Tyr Ile Asp Phe Lys Asp Leu Gly Trp
 355 360 365
 30 His Asp Trp Ile Ile Ala Pro Glu Gly Tyr Gly Ala Phe Tyr Cys Ser
 370 375 380
 Gly Glu Cys Asn Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His
 385 390 395 400
 Ala Ile Val Gln Thr Leu Val His Leu Leu Glu Pro Lys Lys Val Pro
 405 410 415
 40 Lys Pro Cys Cys Ala Pro Thr Arg Leu Gly Ala Leu Pro Val Leu Tyr
 420 425 430
 His Leu Asn Asp Glu Asn Val Asn Leu Lys Lys Tyr Arg Asn Met Ile
 435 440 445
 45 Val Lys Ser Cys Gly Cys His
 450 455

- 134 -

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 104 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10

(ix) FEATURE:

- 15 (A) NAME/KEY: Protein
 (B) LOCATION: 1..104
 (D) OTHER INFORMATION: /note= "BMP3"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

20 Cys Ala Arg Arg Tyr Leu Lys Val Asp Phe Ala Asp Ile Gly Trp Ser
 1 5 10
 Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp Ala Tyr Tyr Cys Ser Gly
 20 25 30
 25 Ala Cys Gln Phe Pro Met Pro Lys Ser Leu Lys Pro Ser Asn His Ala
 35 40 45
 Thr Ile Gln Ser Ile Val Ala Arg Ala Val Gly Val Val Pro Gly Ile
 30 50 55 60
 Pro Glu Pro Cys Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu
 65 70 75 80
 35 Phe Phe Asp Glu Asn Lys Asn Val Val Leu Lys Val Tyr Pro Asn Met
 85 90 95
 Thr Val Glu Ser Cys Ala Cys Arg
 100

40

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 102 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

50

- 135 -

(vi) ORIGINAL SOURCE:

(A) ORGANISM: HOMO SAPIENS

(ix) FEATURE:

5

(A) NAME/KEY: Protein

(B) LOCATION: 1..102

(D) OTHER INFORMATION: /note= "BMP5"

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg Asp Leu Gly Trp Gln
 1 5 10 15

15

Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala Phe Tyr Cys Asp Gly
 20 25 30

Glu Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His Ala
 35 40 45

20

Ile Val Gln Thr Leu Val His Leu Met Phe Pro Asp His Val Pro Lys
 50 55 60

25

Pro Cys Cys Ala Pro Thr Lys Leu Asn Ala Ile Ser Val Leu Tyr Phe
 65 70 75 80

Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr Arg Asn Met Val Val
 85 90 95

30

Arg Ser Cys Gly Cys His
 100

(2) INFORMATION FOR SEQ ID NO:28:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 102 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:

(A) ORGANISM: HOMO SAPIENS

45

(ix) FEATURE:

(A) NAME/KEY: Protein

(B) LOCATION: 1..102

(D) OTHER INFORMATION: /note= "BMP6"

50

- 136 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

5 Cys Arg Lys His Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Gln
 1 5 10 15
 Asp Trp Ile Ile Ala Pro Lys Gly Tyr Ala Ala Asn Tyr Cys Asp Gly
 20 25 30
 10 Glu Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His Ala
 35 40 45
 Ile Val Gln Thr Leu Val His Leu Met Asn Pro Glu Tyr Val Pro Lys
 50 55 60
 15 Pro Cys Cys Ala Pro Thr Lys Leu Asn Ala Ile Ser Val Leu Tyr Phe
 65 70 75 80
 Asp Asp Asn Ser Asn Val Ile Leu Lys Lys Tyr Arg Trp Met Val Val
 85 90 95
 20 Arg Ala Cys Gly Cys His
 100

(2) INFORMATION FOR SEQ ID NO:29:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 102 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

35

(A) NAME/KEY: Protein

(B) LOCATION: 1..102

(D) OTHER INFORMATION: /label= OPX

/note= "WHEREIN EACH XAA IS INDEPENDENTLY SELECTED
 FROM A GROUP OF ONE OR MORE SPECIFIED AMINO ACIDS
 AS DEFINED IN THE SPECIFICATION (SECTION II.B.2.)"

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

45 Cys Xaa Xaa His Glu Leu Tyr Val Xaa Phe Xaa Asp Leu Gly Trp Xaa
 1 5 10 15
 Asp Trp Xaa Ile Ala Pro Xaa Gly Tyr Xaa Ala Tyr Tyr Cys Glu Gly
 20 25 30
 50

- 137 -

[illegible]

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(1x) FEATURE:

- (A) NAME/KEY: Protein
(B) LOCATION: 1..97
(D) OTHER INFORMATION: /label= GENERIC-SEQ5
/note= "WHEREIN EACH XAA IS INDEPENDENTLY SELECTED
FROM A GROUP OF ONE OR MORE SPECIFIED AMINO ACIDS
AS DEFINED IN THE SPECIFICATION."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Leu Xaa Xaa Xaa Phe Xaa Xaa Xaa Gly Trp Xaa Xaa Trp Xaa Xaa Xaa
1 5 10 15

Pro Xaa Xaa Xaa Xaa Ala Xaa Tyr Cys Xaa Gly Xaa Cys Xaa Xaa Pro
 20 25 30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala Xaa Xaa Xaa Xaa Xaa
 35 40 45

- 138 -

[illegible]

(2) INFORMATION FOR SEQ ID NO:31:

```

15      (i) SEQUENCE CHARACTERISTICS:
          (A) LENGTH: 102 amino acids
          (B) TYPE: amino acid
          (C) STRANDEDNESS: single
          (D) TOPOLOGY: linear
20
      (ii) MOLECULE TYPE: protein

      (ix) FEATURE:
          (A) NAME/KEY: Protein
          (B) LOCATION: 1..102
          (D) OTHER INFORMATION: /label= GENERIC-SEQ6
                                  /note= "WHEREIN EACH XAA IS INDEPENDENTLY SELECTED
40                                  FROM A GROUP OF ONE OR MORE SPECIFIED AMINO ACIDS
                                  AS DEFINED IN THE SPECIFICATION. "

      (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

35      Cys Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Phe Xaa Xaa Xaa Gly Trp Xaa
         1           5             10            15
        Xaa Trp Xaa Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala Xaa Tyr Cys Xaa Gly
                                20              25                30
40      Xaa Cys Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala
         35                      40                        45
45      Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
         50                      55                        60

```


- 139 -

Xaa Cys Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa
 65 70 75 80
 Xaa Xaa Xaa Xaa Xaa Val Xaa Leu Xaa Xaa Xaa Xaa Xaa Met Xaa Val
 5 85 90 95
 Xaa Xaa Cys Xaa Cys Xaa
 100

10 (2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1247 base pairs
 (B) TYPE: nucleic acid
 15 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: cDNA
 20 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: HOMO SAPIENS
 (F) TISSUE TYPE: BRAIN
 (ix) FEATURE:
 25 (A) NAME/KEY: CDS
 (B) LOCATION: 84..1199
 (D) OTHER INFORMATION: /product= "GDF-1"
 /note= "GDF-1 CDNA"

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GGGGACACCG GCCCCGCCCT CAGCCCACTG GTCCCGGGCC GCCGCGGACC CTGCGCACTC 60
 35 TCTGGTCATC GCCTGGGAGG AAG ATG CCA CCG CCG CAG CAA GGT CCC TGC 110
 Met Pro Pro Pro Gln Gln Gly Pro Cys
 1 5
 GGC CAC CAC CTC CTC CTC CTC CTG GCC CTG CTG CTG CCC TCG CTG CCC 158
 40 Gly His His Leu Leu Leu Leu Leu Ala Leu Leu Leu Pro Ser Leu Pro 25
 10 15 20 25
 CTG ACC CGC GCC CCC GTG CCC CCA GGC CCA GCC GCC GCC CTG CTC CAG 206
 45 Leu Thr Arg Ala Pro Val Pro Pro Gly Pro Ala Ala Ala Leu Leu Gln 30 35 40
 GCT CTA GGA CTG CGC GAT GAG CCC CAG GGT GCC CCC AGG CTC CGG CCG 254
 50 Ala Leu Gly Leu Arg Asp Glu Pro Gln Gly Ala Pro Arg Leu Arg Pro 45 50 55

- 140 -

	GTT	CCC	CCG	GTC	ATG	TGG	CGC	CTG	TTT	CGA	CGC	CGG	GAC	CCC	CAG	GAG	302
	Val	Pro	Pro	Val	Met	Trp	Arg	Leu	Phe	Arg	Arg	Arg	Asp	Pro	Gln	Glu	
			60					65					70				
5	ACC	AGG	TCT	GGC	TCG	CGG	CGG	ACG	TCC	CCA	GGG	GTC	ACC	CTG	CAA	CCG	350
	Thr	Arg	Ser	Gly	Ser	Arg	Arg	Thr	Ser	Pro	Gly	Val	Thr	Leu	Gln	Pro	
		75					80					85					
10	TGC	CAC	GTG	GAG	GAG	CTG	GGG	GTC	GCC	GGA	AAC	ATC	GTG	CGC	CAC	ATC	398
	Cys	His	Val	Glu	Glu	Leu	Gly	Val	Ala	Gly	Asn	Ile	Val	Arg	His	Ile	
	90					95					100					105	
15	CCG	GAC	CGC	GGT	GCG	CCC	ACC	CGG	GCC	TCG	GAG	CCT	GTC	TCG	GCC	GCG	446
	Pro	Asp	Arg	Gly	Pro	Pro	Thr	Arg	Ala	Ser	Glu	Pro	Val	Ser	Ala	Ala	
					110					115					120		
20	GGG	CAT	TGC	CCT	GAG	TGG	ACA	GTC	GTC	TTC	GAC	CTG	TCG	GCT	GTG	GAA	494
	Gly	His	Cys	Pro	Glu	Trp	Thr	Val	Val	Phe	Asp	Leu	Ser	Ala	Val	Glu	
				125					130					135			
25	CCC	GCT	GAG	CGC	CCG	AGC	CGG	GCC	CGC	CTG	GAG	CTG	CGT	TTC	GCG	GCG	542
	Pro	Ala	Glu	Arg	Pro	Ser	Arg	Ala	Arg	Leu	Glu	Leu	Arg	Phe	Ala	Ala	
			140					145					150				
30	GCG	GCG	GCG	GCA	GCC	CCG	GAG	GGC	GGC	TGG	GAG	CTG	AGC	GTG	GCG	CAA	590
	Ala	Ala	Ala	Ala	Ala	Pro	Glu	Gly	Gly	Trp	Glu	Leu	Ser	Val	Ala	Gln	
			155				160					165					
35	GCG	GGC	CAG	GGC	GCG	GGC	GCG	GAC	CCC	GGG	CCG	GTG	CTG	CTC	CGC	CAG	638
	Ala	Gly	Gln	Gly	Ala	Gly	Ala	Asp	Pro	Gly	Pro	Val	Leu	Leu	Arg	Gln	
	170					175					180					185	
40	TTG	GTG	CCC	GCC	CTG	GGG	CCG	CCA	GTG	CGC	GCG	GAG	CTG	CTG	GGC	GCC	686
	Leu	Val	Pro	Ala	Leu	Gly	Pro	Pro	Val	Arg	Ala	Glu	Leu	Leu	Gly	Ala	
					190					195					200		
45	GCT	TGG	GCT	CGC	AAC	GCC	TCA	TGG	CCG	CGC	AGC	CTC	CGC	CTG	GCG	CTG	734
	Ala	Trp	Ala	Arg	Asn	Ala	Ser	Trp	Pro	Arg	Ser	Leu	Arg	Leu	Ala	Leu	
				205					210					215			
50	GCG	CTA	CGC	CCC	CGG	GCC	CCT	GCC	GCC	TGC	GCG	CGC	CTG	GCC	GAG	GCC	782
	Ala	Leu	Arg	Pro	Arg	Ala	Pro	Ala	Ala	Cys	Ala	Arg	Leu	Ala	Glu	Ala	
			220					225					230				
55	TCG	CTG	CTG	CTG	GTG	ACC	CTC	GAC	CCG	CGC	CTG	TGC	CAC	CCC	CTG	GCC	830
	Ser	Leu	Leu	Leu	Val	Thr	Leu	Asp	Pro	Arg	Leu	Cys	His	Pro	Leu	Ala	
		235					240					245					

- 141 -

	CGG CCG CGG CGC GAC GCC GAA CCC GTG TTG GGC GGC GGC CCC GGG GGC	878
	Arg Pro Arg Arg Asp Ala Glu Pro Val Leu Gly Gly Gly Pro Gly Gly	
	250 255 260 265	
5	GCT TGT CGC GCG CGG CGG CTG TAC GTG AGC TTC CGC GAG GTG GGC TGG	926
	Ala Cys Arg Ala Arg Arg Leu Tyr Val Ser Phe Arg Glu Val Gly Trp	
	270 275 280	
10	CAC CGC TGG GTC ATC GCG CCG CGC GGC TTC CTG GCC AAC TAC TGC CAG	974
	His Arg Trp Val Ile Ala Pro Arg Gly Phe Leu Ala Asn Tyr Cys Gln	
	285 290 295	
15	GGT CAG TGC GCG CTG CCC GTC GCG CTG TCG GGG TCC GGG GGG CCG CCG	1022
	Gly Gln Cys Ala Leu Pro Val Ala Leu Ser Gly Ser Gly Gly Pro Pro	
	300 305 310	
20	GCG CTC AAC CAC GCT GTG CTG CGC GCG CTC ATG CAC GCG GCC GCC CCG	1070
	Ala Leu Asn His Ala Val Leu Arg Ala Leu Met His Ala Ala Ala Pro	
	315 320 325	
25	GGA GCC GCC GAC CTG CCC TGC TGC GTG CCC GCG CGC CTG TCG CCC ATC	1118
	Gly Ala Ala Asp Leu Pro Cys Cys Val Pro Ala Arg Leu Ser Pro Ile	
	330 335 340 345	
30	TCC GTG CTC TTC TTT GAC AAC AGC GAC AAC GTG GTG CTG CGG CAG TAT	1166
	Ser Val Leu Phe Phe Asp Asn Ser Asp Asn Val Val Leu Arg Gln Tyr	
	350 355 360	
35	GAG GAC ATG GTG GTG GAC GAG TGC GGC TGC CGC TAACCCGGGG CGGGCAGGGA	1219
	Glu Asp Met Val Val Asp Glu Cys Gly Cys Arg	
	365 370	
40	CCCGGGCCCA ACAATAAATG CCGCGTGG	1247

35

(2) INFORMATION FOR SEQ ID NO:33:

(1) SEQUENCE CHARACTERISTICS:

40

(A) LENGTH: 372 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Met	Pro	Pro	Pro	Gln	Gln	Gly	Pro	Cys	Gly	His	His	Leu	Leu	Leu	Leu
1				5					10					15	

- 142 -

	Leu	Ala	Leu	Leu	Leu	Pro	Ser	Leu	Pro	Leu	Thr	Arg	Ala	Pro	Val	Pro	
			20						25						30		
5	Pro	Gly	Pro	Ala	Ala	Ala	Leu	Leu	Gln	Ala	Leu	Gly	Leu	Arg	Asp	Glu	
			35					40					45				
	Pro	Gln	Gly	Ala	Pro	Arg	Leu	Arg	Pro	Val	Pro	Pro	Val	Met	Trp	Arg	
		50					55					60					
10	Leu	Phe	Arg	Arg	Arg	Asp	Pro	Gln	Glu	Thr	Arg	Ser	Gly	Ser	Arg	Arg	
	65					70					75					80	
	Thr	Ser	Pro	Gly	Val	Thr	Leu	Gln	Pro	Cys	His	Val	Glu	Glu	Leu	Gly	
					85					90					95		
15	Val	Ala	Gly	Asn	Ile	Val	Arg	His	Ile	Pro	Asp	Arg	Gly	Ala	Pro	Thr	
			100						105					110			
	Arg	Ala	Ser	Glu	Pro	Val	Ser	Ala	Ala	Gly	His	Cys	Pro	Glu	Trp	Thr	
20			115					120					125				
	Val	Val	Phe	Asp	Leu	Ser	Ala	Val	Glu	Pro	Ala	Glu	Arg	Pro	Ser	Arg	
		130					135					140					
25	Ala	Arg	Leu	Glu	Leu	Arg	Phe	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Pro	Glu	
	145					150					155					160	
	Gly	Gly	Trp	Glu	Leu	Ser	Val	Ala	Gln	Ala	Gly	Gln	Gly	Ala	Gly	Ala	
				165					170						175		
30	Asp	Pro	Gly	Pro	Val	Leu	Leu	Arg	Gln	Leu	Val	Pro	Ala	Leu	Gly	Pro	
				180					185					190			
	Pro	Val	Arg	Ala	Glu	Leu	Leu	Gly	Ala	Ala	Trp	Ala	Arg	Asn	Ala	Ser	
35			195					200					205				
	Trp	Pro	Arg	Ser	Leu	Arg	Leu	Ala	Leu	Ala	Leu	Arg	Pro	Arg	Ala	Pro	
		210					215					220					
40	Ala	Ala	Cys	Ala	Arg	Leu	Ala	Glu	Ala	Ser	Leu	Leu	Leu	Val	Thr	Leu	
	225					230					235					240	
	Asp	Pro	Arg	Leu	Cys	His	Pro	Leu	Ala	Arg	Pro	Arg	Arg	Asp	Ala	Glu	
					245					250					255		
45	Pro	Val	Leu	Gly	Gly	Gly	Pro	Gly	Gly	Ala	Cys	Arg	Ala	Arg	Arg	Leu	
			260						265					270			

- 143 -

Tyr Val Ser Phe Arg Glu Val Gly Trp His Arg Trp Val Ile Ala Pro
 275 280 285

5 Arg Gly Phe Leu Ala Asn Tyr Cys Gln Gly Gln Cys Ala Leu Pro Val
 290 295 300

Ala Leu Ser Gly Ser Gly Gly Pro Pro Ala Leu Asn His Ala Val Leu
 305 310 315 320

10 Arg Ala Leu Met His Ala Ala Ala Pro Gly Ala Ala Asp Leu Pro Cys
 325 330 335

Cys Val Pro Ala Arg Leu Ser Pro Ile Ser Val Leu Phe Phe Asp Asn
 340 345 350

15 Ser Asp Asn Val Val Leu Arg Gln Tyr Glu Asp Met Val Val Asp Glu
 355 360 365

20 Cys Gly Cys Arg
 370

- 144 -

What is claimed is:

1. A morphogen-enriched infant formula composition.
- 5 2. The morphogen-enriched infant formula of claim 1 wherein said formula is milk-based.
3. The morphogen-enriched infant formula of claim 1 wherein said formula is soy-based.
- 10 4. The morphogen-enriched infant formula of claim 1 wherein said formula is adapted for preterm infants.
- 15 5. The morphogen-enriched infant formula of claim 1 wherein said formula is adapted for low birth weight infants.
6. A morphogen-enriched dietary composition for individuals at risk for tissue damage due to protein-energy malnutrition.
- 20 7. The morphogen-enriched dietary composition of claim 6 wherein said individual is at risk for tissue damage from starvation, dehydration, anorexia nervosa or trauma.
- 25 8. The morphogen-enriched dietary composition of claim 6 wherein said individual is at risk for tissue damage from a malabsorption-malnutrition disorder.
- 30 9. The morphogen-enriched dietary composition of claim 8 wherein malabsorption-malnutrition disorder results from a digestive or intestinal fistula, short bowel, gastrointestinal disorder or hypercatabolism.

- 145 -

10. The morphogen-enriched dietary composition of claim 6 wherein said individual is at risk for malnutrition-malabsorption induced tissue damage following radiotherapy, chemotherapy or surgery.
- 5
11. A morphogen-enriched dietary composition for individuals at risk for tissue damage due to altered metabolism function.
- 10
12. The morphogen-enriched dietary composition of claim 11 wherein said individual at risk is a pregnant, lactating or post-menopausal female.
- 15
13. The morphogen-enriched dietary composition of claim 11 wherein said individual at risk is an aged individual.
- 20
14. The composition of claim 1, 6 or 11 wherein said morphogen is associated with a controlled release componen, adapted such that the morphogen is released in a controlled manner lower in the gastrointestinal tract.
- 25
15. The composition of claim 1, 6 or 11 adapted for enteral administration.
- 30
16. The composition of claim 1, 6 or 11 adapted for aerosol administration.
17. The composition of claim 1, 6 or 11 adapted for nasal administration.
18. The composition of claim 6 or 11 formulated as a solid.

- 146 -

19. The composition of claim 18 wherein said solid is a tablet, troche or lozenge.
- 5 20. The composition of claim 1, 6 or 11 formulated as a liquid.
21. The composition of claim 20 wherein said liquid is a beverage or a syrup.
- 10 22. The composition of claim 1, 6 or 11 wherein said morphogen is associated with a morphogen-solubilizing molecule.
- 15 23. The composition of claim 22 wherein said molecule is casein or a derivative, salt or analog thereof.
24. The composition of claim 22 wherein said molecule comprises part or all of a morphogen pro domain.
- 20 25. The composition of claim 1, 6 or 11 wherein said morphogen comprises an amino acid sequence sharing at least 70% homology with one of the sequences selected from the group consisting of: OP-1, OP-2, CBMP2, BMP3(fx), Vgl(fx), Vgr(fx), DPP(fx), GDF-1(fx) and 25 60A(fx).
26. The composition of claim 25 wherein said morphogen comprises an amino acid sequence sharing a last 80% homology with one of the sequences selected from the group consisting of: OP-1, OP-2, CBMP2, BMP3(fx), 30 BMP5(fx), BMP6(fx), Vgl(fx), Vgr(fx), DPP(fx), GDF-1(fx) and 60A(fx).

- 147 -

27. The composition of claim 1, 6 or 11 wherein said morphogen comprises an amino acid sequence having greater than 60% amino acid identity with the sequence defined by residues 43-139 of Seq. ID No. 5 (hOP1).
- 5
28. The composition of claim 27 wherein said morphogen comprises an amino acid sequence having greater than 65% amino acid identity with the sequence defined by residues 43-139 of Seq. ID No. 5 (hOP1).
- 10
29. The composition of claim 28 wherein said morphogen comprises an amino acid sequence defined by residues 43-139 of Seq. ID No. 5 (hOP1), including allelic and species variants thereof.
- 15
30. The composition of claim 1, 6 or 11 wherein said morphogen comprises an amino acid sequence defined by Generic Sequences 1, 2, 3, 4, 5 or 6 (Seq. ID Nos. 1, 2, 3, 4, 30 or 31).
- 20
31. The composition of claim 1, 6 or 11 wherein said morphogen comprises an amino acid sequence defined by OPX (Seq. ID No. 29).
- 25
32. The composition of claim 1, 6 or 11 wherein said morphogen comprises an amino acid sequence encoded by a nucleic acid that hybridizes under stringent conditions with a DNA defined by nucleotides 1036-1341 of Seq. ID No. 16 or nucleotides 1390-1695 of Seq. ID No. 20.
- 30

- 148 -

- 5 33. A method for improving the human milk mimetic characteristics of an infant formula, the method comprising the step of adding a morphogenically effective concentration of a morphogen to said formula prior to providing said formula to an infant.
- 10 34. A method for enhancing tissue viability in a malnourished individual, the method comprising the step of providing to said individual a morphogenically effective concentration of a morphogen.
- 15 35. A method for enhancing tissue viability in an individual having altered metabolic function, the method comprising the step of providing to said individual a morphogenically effective concentration of a morphogen.
- 20 36. The method of claim 35 wherein said individual is a pregnant, lactating or postmenopausal female.
- 25 37. The method of claim 35 wherein said individual is an aged individual.
38. The method of claim 34 or 35 wherein said morphogen is provided to said individual as part of a food formulation.
39. The method of claim 34 or 35 wherein said morphogen is provided to said individual as a dietary supplement.

- 149 -



40. The method of claim 33, 34 or 35 wherein said morphogen comprises an amino acid sequence sharing at least 70% homology with one of the sequences selected from the group consisting of: OP-1, OP-2, CBMP2, BMP3(fx), Vg1(fx), Vgr(fx), DPP(fx), GDF-1(fx) and 60A(fx).
41. The method of claim 40 wherein said morphogen comprises an amino acid sequence sharing a last 80% homology with one of the sequences selected from the group consisting of: OP-1, OP-2, CBMP2, BMP3(fx), BMP5(fx), BMP6(fx), Vg1(fx), Vgr(fx), DPP(fx), GDF-1(fx) and 60A(fx).
42. The method of claim 33, 34 or 35 wherein said morphogen comprises an amino acid sequence having greater than 60% amino acid identity with the sequence defined by residues 43-139 of Seq. ID No. 5 (hOP1).
43. The method of claim 42 wherein said morphogen comprises an amino acid sequence having greater than 65% amino acid identity with the sequence defined by residues 43-139 of Seq. ID No. 5 (hOP1).
44. The method of claim 43 wherein said morphogen comprises an amino acid sequence defined by residues 43-139 of Seq. ID No. 5 (hOP1), including allelic and species variants thereof.
45. The method of claim 33, 34 or 35 wherein said morphogen comprises an amino acid sequence defined by Generic Sequences 1, 2, 3, 4, 5 or 6 (Seq. ID Nos. 1, 2, 3, 4, 30 or 31).

- 150 -

46. The method of claim 33, 34 or 35 wherein said morphogen comprises an amino acid sequence defined by OPX (Seq. ID No. 29).
- 5 47. The method of claim 33, 34 or 35 wherein said morphogen comprises an amino acid sequence encoded by a nucleic acid that hybridizes under stringent conditions with a DNA defined by nucleotides 1036-1341 of Seq. ID No. 16 or nucleotides 1390-1695 of Seq. ID No. 20.
- 10
48. The invention of claim 1, 6, 11, 33, 34 or 35 wherein said morphogen comprises a dimeric protein species complexed with a peptide comprising a pro region of a member of the morphogen family, or an allelic, species or other sequence variant thereof.
- 15
49. The invention of claim 48 wherein said dimeric morphogen species is noncovalently complexed with said peptide.
- 20 50. The invention of claim 48 or 49 wherein said dimeric morphogen species is complexed with two said peptides.
51. The invention of claim 48 or 49 wherein said peptide comprises at least the first eighteen amino acids of a sequence defining said pro region.
- 25
52. The invention of claim 51 wherein said peptide comprises the full length of said pro region.

- 151 -

53. The invention of claim 48 or 49 wherein said peptide comprises a nucleic acid that hybridizes under stringent conditions with a DNA defined by nucleotides 136-192 of Seq. ID No. 16, or nucleotides 157-211 of Seq. ID No. 20.
54. The invention of claim 48 or 49 wherein said complex is further stabilized by exposure to a basic amino acid, a detergent or a carrier protein.

*Fig. 1A**Fig. 1B*




Fig. 2A




Fig. 2B

3/8




Fig. 3A








Fig. 3B

4/8

*Fig. 4A**Fig. 4B*

SUBSTITUTE SHEET

*Fig. 5**Fig. 7*

SUBSTITUTE SHEET

6/8




Fig. 6A




Fig. 6B

7/8





Fig. 8A

*Fig. 8B*